10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CD152-AcalephFluor568-labled

Catalog # AP101318

Product Information

Application	FC, IF/IC
Primary Accession	P16410
Reactivity	Human
Clonality	Monoclonal
Calculated MW	24656

Additional Information

Gene ID	1493
Other Names	CD152; Cytotoxic T-lymphocyte protein 4; Cytotoxic T-lymphocyte-associated antigen 4; CTLA-4; CD antigen CD152
Target/Specificity	Recognizes human CD152
Dilution	FC~~1:10~50 IF/IC~~N/A
Format	Mouse IgG1 kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	CTLA4
Synonyms	CD152
Function	Inhibitory receptor acting as a major negative regulator of T-cell responses (PubMed:11279501, PubMed:11279502, PubMed:16551244, PubMed:1714933, PubMed:18641304, PubMed:28484017). Acts as a decoy receptor: the affinity of CTLA4 for its natural B7 family ligands, CD80 and CD86, is considerably stronger than the affinity of their cognate stimulatory coreceptor CD28 (PubMed:11279501, PubMed:11279502, PubMed:16551244, PubMed:1714933, PubMed:28484017).
Cellular Location	Cell membrane; Single-pass type I membrane protein. Note=Exists primarily an intracellular antigen whose surface expression is tightly regulated by restricted trafficking to the cell surface and rapid internalization
Tissue Location	Widely expressed with highest levels in lymphoid tissues. Detected in activated T-cells where expression levels are 30- to 50-fold less than CD28, the stimulatory coreceptor, on the cell surface following activation.

Background

AcalephFluor568-labled mouse monoclonal antibody to CD152

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.