10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CD33-AcalephFluor647-labled

Catalog # AP101446

Product Information

Application	FC, IF/IC
Primary Accession	P20138
Reactivity	Human
Clonality	Monoclonal
Calculated MW	39825

Additional Information

Gene ID	945
Other Names	SIGLEC3; Myeloid cell surface antigen CD33; Sialic acid-binding Ig-like lectin 3; Siglec-3; gp67; CD antigen CD33
Target/Specificity	Recognizes human CD33
Dilution	FC~~1:10~50 IF/IC~~N/A
Format	Mouse IgG1. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	CD33
Synonyms	SIGLEC3
Function	Sialic-acid-binding immunoglobulin-like lectin (Siglec) that plays a role in mediating cell-cell interactions and in maintaining immune cells in a resting state (PubMed:10611343, PubMed:11320212, PubMed:15597323). Preferentially recognizes and binds alpha-2,3- and more avidly alpha-2,6-linked sialic acid-bearing glycans (PubMed:7718872). Upon engagement of ligands such as C1q or syalylated glycoproteins, two immunoreceptor tyrosine-based inhibitory motifs (ITIMs) located in CD33 cytoplasmic tail are phosphorylated by Src-like kinases such as LCK (PubMed:10887109, PubMed:28325905). These phosphorylations provide docking sites for the recruitment and activation of protein-tyrosine phosphatases PTPN6/SHP-1 and PTPN11/SHP- 2 (PubMed:10206955, PubMed:10556798, PubMed:10887109). In turn, these phosphatases regulate downstream pathways through dephosphorylation of signaling molecules (PubMed:10206955, PubMed:10887109). One of the repressive effect of CD33 on monocyte activation requires phosphoinositide 3-kinase/PI3K (PubMed:15597323).

Cellular Location	[Isoform CD33M]: Cell membrane; Single-pass type I membrane protein
Tissue Location	Monocytic/myeloid lineage cells. In the brain, CD33 is mainly expressed on microglial cells

Background

AcalephFluor647-labled mouse monoclonal antibody to CD33

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.