

# SLC3A2 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP10152b

## Product Information

---

<b>Application</b>	WB, IHC-P, E
<b>Primary Accession</b>	<a href="#">P08195</a>
<b>Other Accession</b>	<a href="#">NP_001012681.1</a> , <a href="#">NP_001012679.1</a> , <a href="#">NP_002385.3</a> , <a href="#">NP_001012680.1</a> , <a href="#">NP_001013269.1</a> , <a href="#">NP_001012682.1</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB22529
<b>Calculated MW</b>	67994
<b>Antigen Region</b>	588-615

## Additional Information

---

<b>Gene ID</b>	6520
<b>Other Names</b>	4F2 cell-surface antigen heavy chain, 4F2hc, 4F2 heavy chain antigen, Lymphocyte activation antigen 4F2 large subunit, Solute carrier family 3 member 2, CD98, SLC3A2, MDU1
<b>Target/Specificity</b>	This SLC3A2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 588-615 amino acids from the C-terminal region of human SLC3A2.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	SLC3A2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.


## Protein Information

---


<b>Name</b>	SLC3A2 ( <a href="#">HGNC:11026</a> )
<b>Synonyms</b>	MDU1

<b>Function</b>	<p>Acts as a chaperone that facilitates biogenesis and trafficking of functional transporters heterodimers to the plasma membrane. Forms heterodimer with SLC7 family transporters (SLC7A5, SLC7A6, SLC7A7, SLC7A8, SLC7A10 and SLC7A11), a group of amino-acid antiporters (PubMed:<a href="#">10574970</a>, PubMed:<a href="#">10903140</a>, PubMed:<a href="#">11557028</a>, PubMed:<a href="#">30867591</a>, PubMed:<a href="#">33298890</a>, PubMed:<a href="#">33758168</a>, PubMed:<a href="#">34880232</a>, PubMed:<a href="#">9751058</a>, PubMed:<a href="#">9829974</a>, PubMed:<a href="#">9878049</a>). Heterodimers function as amino acids exchangers, the specificity of the substrate depending on the SLC7A subunit. Heterodimers SLC3A2/SLC7A6 or SLC3A2/SLC7A7 mediate the uptake of dibasic amino acids (PubMed:<a href="#">10903140</a>, PubMed:<a href="#">9829974</a>). Heterodimer SLC3A2/SLC7A11 functions as an antiporter by mediating the exchange of extracellular anionic L-cystine and intracellular L-glutamate across the cellular plasma membrane (PubMed:<a href="#">34880232</a>). SLC3A2/SLC7A10 translocates small neutral L- and D- amino acids across the plasma membrane (By similarity). SLC3A2/SLC75 or SLC3A2/SLC7A8 translocates neutral amino acids with broad specificity, thyroid hormones and L-DOPA (PubMed:<a href="#">10574970</a>, PubMed:<a href="#">11389679</a>, PubMed:<a href="#">11557028</a>, PubMed:<a href="#">11564694</a>, PubMed:<a href="#">11742812</a>, PubMed:<a href="#">12117417</a>, PubMed:<a href="#">12225859</a>, PubMed:<a href="#">12716892</a>, PubMed:<a href="#">15980244</a>, PubMed:<a href="#">30867591</a>, PubMed:<a href="#">33298890</a>, PubMed:<a href="#">33758168</a>). SLC3A2 is essential for plasma membrane localization, stability, and the transport activity of SLC7A5 and SLC7A8 (PubMed:<a href="#">10391915</a>, PubMed:<a href="#">10574970</a>, PubMed:<a href="#">11311135</a>, PubMed:<a href="#">15769744</a>, PubMed:<a href="#">33066406</a>). When associated with LAPTM4B, the heterodimer SLC7A5 is recruited to lysosomes to promote leucine uptake into these organelles, and thereby mediates mTORC1 activation (PubMed:<a href="#">25998567</a>). Modulates integrin-related signaling and is essential for integrin-dependent cell spreading, migration and tumor progression (PubMed:<a href="#">11121428</a>, PubMed:<a href="#">15625115</a>).</p>
<b>Cellular Location</b>	<p>Apical cell membrane. Cell membrane; Single-pass type II membrane protein. Cell junction {ECO:0000250 UniProtKB:P10852}. Lysosome membrane. Melanosome. Basolateral cell membrane {ECO:0000250 UniProtKB:P10852}. Note=Localized at the plasma membrane when associated with SLC7A5/LAT1 or SLC7A8/LAT2 (PubMed:11311135, PubMed:9751058). Localized to the apical membrane of placental syncytiotrophoblastic cells (PubMed:11742812). Recruited to lysosomes by LAPTM4B (PubMed:25998567). Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065) Located selectively at cell-cell adhesion sites (By similarity) Colocalized with SLC7A8/LAT2 at the basolateral membrane of kidney proximal tubules and small intestine epithelia. Expressed in both luminal and abluminal membranes of brain capillary endothelial cells (By similarity). {ECO:0000250 UniProtKB:P10852, ECO:0000269 PubMed:11311135, ECO:0000269 PubMed:11742812, ECO:0000269 PubMed:17081065, ECO:0000269 PubMed:25998567, ECO:0000269 PubMed:9751058}</p>
<b>Tissue Location</b>	<p>Expressed ubiquitously in all tissues tested with highest levels detected in kidney, placenta and testis and weakest level in thymus. During gestation, expression in the placenta was significantly stronger at full-term than at the mid-trimester stage Expressed in HUVECS and at low levels in resting peripheral blood T- lymphocytes and quiescent fibroblasts. Also expressed in fetal liver and in the astrocytic process of primary astrocytic gliomas. Expressed in retinal endothelial cells and in the intestinal epithelial cell line C2BBel.</p>


## Images




sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



AP10152b staining SLC3A2 in human placenta tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



All lanes : Anti-SLC3A2 Antibody (C-term) at 1:4000 dilution Lane 1: HL-60 whole cell lysate Lane 2: HepG2 whole cell lysate Lane 3: A431 whole cell lysate Lane 4: Human kidney lysate Lane 5: Human placenta lysate Lane 6: Hela whole cell lysate Lysates/proteins at 10 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 68 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-SLC3A2 Antibody (C-term) at 1:2000 dilution + human kidney lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 68 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.