10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CD50-AcalephFluor488-labled

Catalog # AP101892

Product Information

Application	FC, IF/IC
Primary Accession	P32942
Reactivity	Human
Clonality	Monoclonal
Calculated MW	59541

Additional Information

Gene ID	3385
Other Names	Intercellular adhesion molecule 3; ICAM-3; CDw50; ICAM-R; CD50
Target/Specificity	Recognizes human CD50
Dilution	FC~~1:10~50 IF/IC~~N/A
Format	Mouse IgG2b kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	ICAM3 (HGNC:5346)
Function	Cell adhesion molecule that functions as a receptor ligand of the signaling receptor ITGAL:ITGB2/LFA-1 (lymphocyte-function associated (LFA) molecule 1) ensuring apoptotic neutrophil phagocytosis by macrophages (PubMed:1448173, PubMed:1448174, PubMed:23775590, PubMed:8325327, PubMed:8459213). Is also a ligand for integrin ITGAD/ITGB2 (PubMed:8777714). Ligand of CD209 on dendritic cells (DC) through an integrin-independent mechanism that requires Ca(2+) and that forms a first contact between DC and resting T cell, facilitating the downstream DC-T cell clustering process and DC-induced proliferation of resting T Cells (PubMed:10721994).
Cellular Location	Cell membrane; Single-pass type I membrane protein Note=Localization at uropodium is mediated through interaction with MSN.
Tissue Location	Constitutively expressed only on resting leukocytes (PubMed:8325327). No expressed on resting or cytokine-activated endothelial cells (PubMed:1448174).

Background

AcalephFluor488-labled mouse monoclonal antibody to CD50

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.