

NOS1AP Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP10269c

Product Information

Application	FC, WB, E
Primary Accession	O75052
Other Accession	O54960 , Q9D3A8 , NP_055512.1 , NP_001158229.1
Reactivity	Human
Predicted	Mouse, Rat, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB27629
Calculated MW	56150
Antigen Region	437-466

Additional Information

Gene ID	9722
Other Names	Carboxyl-terminal PDZ ligand of neuronal nitric oxide synthase protein, C-terminal PDZ ligand of neuronal nitric oxide synthase protein, Nitric oxide synthase 1 adaptor protein, NOS1AP, CAPON, KIAA0464
Target/Specificity	This NOS1AP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 437-466 amino acids from the Central region of human NOS1AP.
Dilution	FC~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	NOS1AP Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NOS1AP (HGNC:16859)
Synonyms	CAPON, KIAA0464

Function	Adapter protein involved in neuronal nitric-oxide (NO) synthesis regulation via its association with nNOS/NOS1. The complex formed with NOS1 and synapsins is necessary for specific NO and synapsin functions at a presynaptic level. Mediates an indirect interaction between NOS1 and RASD1 leading to enhance the ability of NOS1 to activate RASD1. Competes with DLG4 for interaction with NOS1, possibly affecting NOS1 activity by regulating the interaction between NOS1 and DLG4 (By similarity). In kidney podocytes, plays a role in podosomes and filopodia formation through CDC42 activation (PubMed: 33523862).
Cellular Location	Cell projection, filopodium {ECO:0000250 UniProtKB:O54960}. Cell projection, podosome {ECO:0000250 UniProtKB:O54960}
Tissue Location	Expressed in kidney glomeruli podocytes.


Background

This gene encodes a cytosolic protein that binds to the signaling molecule, neuronal nitric oxide synthase (nNOS). This protein has a C-terminal PDZ-binding domain that mediates interactions with nNOS and an N-terminal phosphotyrosine binding (PTB) domain that binds to the small monomeric G protein, Dexas1. Studies of the related mouse and rat proteins have shown that this protein functions as an adapter protein linking nNOS to specific targets, such as Dexas1 and the synapsins. Alternative splicing results in multiple transcript variants encoding different isoforms.

References


Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
 Lu, J., et al. Diabet. Med. 27(9):1074-1079(2010)
 Husted, J.A., et al. Schizophr. Res. 121 (1-3), 187-192 (2010) :
 Tomas, M., et al. J. Am. Coll. Cardiol. 55(24):2745-2752(2010)
 Delorme, R., et al. BMC Med. Genet. 11, 108 (2010) :

Images



Western blot analysis of lysate from MCF-7 cell line, using NOS1AP Antibody (Center)(Cat. #AP10269c). AP10269c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug per lane.

NOS1AP Antibody (Center) (Cat. #AP10269c) flow cytometric analysis of HL-60 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.