

C11orf63 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP10323b

Product Information

Application	WB, FC, E
Primary Accession	Q6NUN7
Other Accession	Q2T9M9 , NP_954575.1 , NP_079082.2
Reactivity	Human, Rat, Mouse
Predicted	Bovine, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB27866
Calculated MW	88569
Antigen Region	659-688

Additional Information

Gene ID	79864
Other Names	Uncharacterized protein C11orf63, C11orf63
Target/Specificity	This C11orf63 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 659-688 amino acids from the C-terminal region of human C11orf63.
Dilution	WB~~1:1000 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	C11orf63 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	JHY (HGNC:26288)
Synonyms	C11orf63
Function	Required for the normal development of cilia in brain ependymal cells lining

the ventricular surfaces.


References

Rose, J. Phd, et al. Mol. Med. (2010) In press :


Images



All lanes : Anti-C11orf63 Antibody (C-term) at 1:1000 dilution
Lane 1: mouse brain tissue lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size : 87kDa
Blocking/Dilution buffer: 5% NFDM/TBST.



C11orf63 Antibody (C-term) (Cat. #AP10323b) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the C11orf63 antibody detected the C11orf63 protein (arrow).



C11orf63 Antibody (C-term) (Cat. #AP10323b) flow cytometric analysis of CEM cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.