

KEAP1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP10441b

Product Information

Application	WB, FC, E
Primary Accession	Q14145
Other Accession	Q684M4 , NP_036421.2 , NP_987096.1
Reactivity	Human
Predicted	Pig
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB28438
Calculated MW	69666
Antigen Region	429-459aa

Additional Information

Gene ID	9817
Other Names	Kelch-like ECH-associated protein 1, Cytosolic inhibitor of Nrf2, INrf2, Kelch-like protein 19, KEAP1, INRF2, KIAA0132, KLHL19
Target/Specificity	This KEAP1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the C-terminal region of human KEAP1.
Dilution	WB~~1:1000 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	KEAP1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	KEAP1 {ECO:0000303 PubMed:14585973, ECO:0000312 HGNC:HGNC:23177}
Function	Substrate-specific adapter of a BCR (BTB-CUL3-RBX1) E3 ubiquitin ligase

complex that regulates the response to oxidative stress by targeting NFE2L2/NRF2 for ubiquitination (PubMed:[14585973](#), PubMed:[15379550](#), PubMed:[15572695](#), PubMed:[15601839](#), PubMed:[15983046](#), PubMed:[37339955](#)). KEAP1 acts as a key sensor of oxidative and electrophilic stress: in normal conditions, the BCR(KEAP1) complex mediates ubiquitination and degradation of NFE2L2/NRF2, a transcription factor regulating expression of many cytoprotective genes (PubMed:[15601839](#), PubMed:[16006525](#)). In response to oxidative stress, different electrophile metabolites trigger non-enzymatic covalent modifications of highly reactive cysteine residues in KEAP1, leading to inactivate the ubiquitin ligase activity of the BCR(KEAP1) complex, promoting NFE2L2/NRF2 nuclear accumulation and expression of phase II detoxifying enzymes (PubMed:[16006525](#), PubMed:[17127771](#), PubMed:[18251510](#), PubMed:[19489739](#), PubMed:[29590092](#)). In response to selective autophagy, KEAP1 is sequestered in inclusion bodies following its interaction with SQSTM1/p62, leading to inactivation of the BCR(KEAP1) complex and activation of NFE2L2/NRF2 (PubMed:[20452972](#)). The BCR(KEAP1) complex also mediates ubiquitination of SQSTM1/p62, increasing SQSTM1/p62 sequestering activity and degradation (PubMed:[28380357](#)). The BCR(KEAP1) complex also targets BPTF and PGAM5 for ubiquitination and degradation by the proteasome (PubMed:[15379550](#), PubMed:[17046835](#)).


Cellular Location

Cytoplasm. Nucleus. Note=Mainly cytoplasmic (PubMed:15601839). In response to selective autophagy, relocalizes to inclusion bodies following interaction with SQSTM1/p62 (PubMed:20452972).


Tissue Location

Broadly expressed, with highest levels in skeletal muscle.

Images



All lanes: Anti-KEAP1 Antibody (C-term) at 1:2000 dilution
 Lane 1: NIH/3T3 whole cell lysate Lane 2: K562 whole cell lysate Lane 3: HepG2 whole cell lysate Lane 4: Jurkat whole cell lysate Lane 5: MCF-7 whole cell lysate
 Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 70 KDa
 Blocking/Dilution buffer: 5% NFDM/TBST.



KEAP1 Antibody (C-term) (Cat. #AP10441b) flow cytometric analysis of HepG2 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.