

# PAX7 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP10488B

### **Product Information**

**Application** IHC-P, WB, E **Primary Accession** P23759

Other Accession P47239, NP 001128726.1, NP 039236.1, NP 002575.1

**Reactivity** Human, Mouse

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Clone Names RB28560
Calculated MW 55119
Antigen Region 410-437

# **Additional Information**

**Gene ID** 5081

Other Names Paired box protein Pax-7, HuP1, PAX7, HUP1

Target/Specificity This PAX7 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 410-437 amino acids from the

C-terminal region of human PAX7.

**Dilution** IHC-P~~1:100~500 WB~~1:1000 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** PAX7 Antibody (C-term) is for research use only and not for use in diagnostic

or therapeutic procedures.

#### **Protein Information**

Name PAX7

Synonyms HUP1

**Function** Transcription factor that is involved in the regulation of muscle stem cells

proliferation, playing a role in myogenesis and muscle regeneration.


# **Background**

PAX7 is a member of the paired box (PAX) family of transcription factors. Members of this gene family typically contain a paired box domain, an octapeptide, and a paired-type homeodomain. These genes play critical roles during fetal development and cancer growth. The specific function of the paired box 7 gene is unknown but speculated to involve tumor suppression since fusion of this gene with a forkhead domain family member has been associated with alveolar rhabdomyosarcoma. Three transcript variants encoding different isoforms have been found for this gene.

## References

Shimada, M., et al. Hum. Genet. 128(4):433-441(2010) Williamson, D., et al. J. Clin. Oncol. 28(13):2151-2158(2010) Xynos, A., et al. Stem Cells 28(5):965-973(2010) Edwards, T.L., et al. Ann. Hum. Genet. 74(2):97-109(2010) Yerges, L.M., et al. J. Bone Miner. Res. 24(12):2039-2049(2009)

# **Images**



Anti-PAX7 Antibody (C-term) at 1:1000 dilution + Mouse skeletal muscle lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 55 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.