

# MICALL1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP10498a

## **Product Information**

**Application** WB, IHC-P, E **Primary Accession Q8N3F8** Other Accession NP 203744.1 Reactivity Human Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB24168 Calculated MW 93441 752-781 **Antigen Region** 

# **Additional Information**

**Gene ID** 85377

Other Names MICAL-like protein 1, Molecule interacting with Rab13, MIRab13, MICALL1,

**KIAA1668, MIRAB13** 

Target/Specificity This MICALL1 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 752-781 amino acids from the

C-terminal region of human MICALL1.

**Dilution** WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** MICALL1 Antibody (C-term) is for research use only and not for use in

diagnostic or therapeutic procedures.

### **Protein Information**

Name MICALL1

Synonyms KIAA1668, MIRAB13

**Function** Lipid-binding protein with higher affinity for phosphatidic acid, a lipid

enriched in recycling endosome membranes. On endosome membranes, acts as a downstream effector of Rab proteins recruiting cytosolic proteins to regulate membrane tubulation (PubMed:19864458, PubMed:20801876, PubMed:23596323, PubMed:34100897). Involved in a late step of receptor-mediated endocytosis regulating for instance endocytosed-EGF receptor trafficking (PubMed:21795389). Alternatively, regulates slow endocytic recycling of endocytosed proteins back to the plasma membrane (PubMed:19864458). Also involved in cargo protein delivery to the plasma membrane (PubMed:34100897). Plays a role in ciliogenesis coordination, recruits EHD1 to primary cilium where it is anchored to the centriole through interaction with tubulins (PubMed:31615969). May indirectly play a role in neurite outgrowth (By similarity).

#### **Cellular Location**

Recycling endosome membrane; Peripheral membrane protein. Late endosome membrane. Cell projection, cilium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole Note=Localization to late endosomes is actin-dependent. Association to tubular recycling endosomes is regulated by RAB35 and ARF6 (PubMed:21951725). Interaction with tubulins achors MICALL1 to the centriole (PubMed:31615969).


# **Background**

MICALL1 may be a cytoskeletal regulator.

## References


Sharma, M., et al. Mol. Biol. Cell 20(24):5181-5194(2009) Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007) Olsen, J.V., et al. Cell 127(3):635-648(2006) Jin, J., et al. Curr. Biol. 14(16):1436-1450(2004) Beausoleil, S.A., et al. Proc. Natl. Acad. Sci. U.S.A. 101(33):12130-12135(2004)

# **Images**



MICALL1 Antibody (C-term) (Cat. #AP10498a) western blot analysis in U251 cell line lysates (35ug/lane). This demonstrates the MICALL1 antibody detected the LACRT protein (arrow).

MICALL1 Antibody (C-term) (Cat. #AP10498a) immunohistochemistry analysis in formalin fixed and paraffin embedded human prostate carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the MICALL1 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.