

ZNF98 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10523b

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	A6NK75
Other Accession	Q9P255
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB28157
Calculated MW	65799
Antigen Region	520-549

Additional Information

Gene ID	148198
Other Names	Zinc finger protein 98, Zinc finger protein 739, Zinc finger protein F7175, ZNF98, ZNF739
Target/Specificity	This ZNF98 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 520-549 amino acids from the C-terminal region of human ZNF98.
Dilution	WB~1:1000 IHC-P~1:100~500 FC~1:10~50 E~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ZNF98 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	ZNF98
Synonyms	ZNF739

Function	May be involved in transcriptional regulation.
Cellular Location	Nucleus.


Background

May be involved in transcriptional regulation.


Images



ZNF98 Antibody (C-term) (Cat. #AP10523b) western blot analysis in MDA-MB435 cell line lysates (35ug/lane). This demonstrates the ZNF98 antibody detected the ZNF98 protein (arrow).



ZNF98 antibody (C-term) (Cat. #AP10523b) immunohistochemistry analysis in formalin fixed and paraffin embedded human lymphnode followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the ZNF98 antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



ZNF98 Antibody (C-term) (Cat. #AP10523b) flow cytometric analysis of MDA-MB435 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.