

# ZNF667 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP10552c

## Product Information

---

Application	WB, IHC-P, FC, E
Primary Accession	<a href="#">Q5HYK9</a>
Other Accession	<a href="#">NP_071386.3</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB28195
Calculated MW	70161
Antigen Region	303-331

## Additional Information

---

Gene ID	63934
Other Names	Zinc finger protein 667, ZNF667
Target/Specificity	This ZNF667 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 303-331 amino acids from the Central region of human ZNF667.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ZNF667 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information


---

Name	ZNF667
Function	May be involved in transcriptional regulation.
Cellular Location	Nucleus.


## Background

May be involved in transcriptional regulation (By similarity).


## Images



ZNF667 Antibody (Center) (Cat. #AP10552c) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the ZNF667 antibody detected the ZNF667 protein (arrow).



ZNF667 Antibody (Center) (Cat. #AP10552c) immunohistochemistry analysis in formalin fixed and paraffin embedded human lung carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the ZNF667 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



ZNF667 Antibody (Center) (Cat. #AP10552c) flow cytometric analysis of CEM cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.