

WDR21C Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP10619a

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	Q8NA75
Other Accession	NP_689631.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB28363
Calculated MW	43747
Antigen Region	1-30

Additional Information

Gene ID	138009
Other Names	DDB1- and CUL4-associated factor 4-like protein 2, WD repeat-containing protein 21C, DCAF4L2, WDR21C
Target/Specificity	This WDR21C antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human WDR21C.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	WDR21C Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Name	DCAF4L2
Synonyms	WDR21C


References

Gerhard, D.S., et al. Genome Res. 14 (10B), 2121-2127 (2004) :


Images



WDR21C Antibody (N-term) (Cat. #AP10619a) western blot analysis in A2058 cell line lysates (35ug/lane). This demonstrates the WDR21C antibody detected the WDR21C protein (arrow).



WDR21C antibody (N-term) (Cat. #AP10619a) immunohistochemistry analysis in formalin fixed and paraffin embedded human colon carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the WDR21C antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



WDR21C Antibody (N-term) (Cat. #AP10619a) flow cytometric analysis of A2058 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.