

# CD11b Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP10668a

## Product Information

---

<b>Application</b>	FC, WB, IHC-P, E
<b>Primary Accession</b>	<a href="#">P11215</a>
<b>Other Accession</b>	<a href="#">NP_000623</a>
<b>Reactivity</b>	Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB17991
<b>Calculated MW</b>	127179
<b>Antigen Region</b>	253-282

## Additional Information

---

<b>Gene ID</b>	3684
<b>Other Names</b>	Integrin alpha-M, CD11 antigen-like family member B, CR-3 alpha chain, Cell surface glycoprotein MAC-1 subunit alpha, Leukocyte adhesion receptor MO1, Neutrophil adherence receptor, CD11b, ITGAM, CD11B, CR3A
<b>Target/Specificity</b>	This CD11b antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 253-282 amino acids from the N-terminal region of human CD11b.
<b>Dilution</b>	FC~~1:10~50 WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	CD11b Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	ITGAM
<b>Synonyms</b>	CD11B, CR3A

<b>Function</b>	Integrin ITGAM/ITGB2 is implicated in various adhesive interactions of monocytes, macrophages and granulocytes as well as in mediating the uptake of complement-coated particles and pathogens (PubMed: <a href="#">20008295</a> , PubMed: <a href="#">9558116</a> ). It is identical with CR-3, the receptor for the iC3b fragment of the third complement component. It probably recognizes the R-G-D peptide in C3b. Integrin ITGAM/ITGB2 is also a receptor for fibrinogen, factor X and ICAM1. It recognizes P1 and P2 peptides of fibrinogen gamma chain. Regulates neutrophil migration (PubMed: <a href="#">28807980</a> ). In association with beta subunit ITGB2/CD18, required for CD177-PRTN3-mediated activation of TNF primed neutrophils (PubMed: <a href="#">21193407</a> ). May regulate phagocytosis-induced apoptosis in extravasated neutrophils (By similarity). May play a role in mast cell development (By similarity). Required with TYROBP/DAP12 in microglia to control production of microglial superoxide ions which promote the neuronal apoptosis that occurs during brain development (By similarity).
<b>Cellular Location</b>	Cell membrane; Single-pass type I membrane protein. Membrane raft; Single-pass type I membrane protein
<b>Tissue Location</b>	Predominantly expressed in monocytes and granulocytes (PubMed:1346576). Expressed in neutrophils (at protein level) (PubMed:21193407).


## Background

CD11b is the integrin alpha M chain. Integrins are heterodimeric integral membrane proteins composed of an alpha chain and a beta chain. This I-domain containing alpha integrin combines with the beta 2 chain (ITGB2) to form a leukocyte-specific integrin referred to as macrophage receptor 1 ('Mac-1'), or inactivated-C3b (iC3b) receptor 3 ('CR3'). The alpha M beta 2 integrin is important in the adherence of neutrophils and monocytes to stimulated endothelium, and also in the phagocytosis of complement coated particles.

## References


Gjelstrup, L.C., et al. J. Immunol. 185(7):4154-4168(2010)  
 Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)  
 Warchol, T., et al. DNA Cell Biol. (2010) In press :  
 Fan, Y., et al. J Eur Acad Dermatol Venereol (2010) In press :  
 Pliyev, B.K., et al. Biochem. Biophys. Res. Commun. 397(2):277-282(2010)

## Images




Anti-CD11b Antibody (N-term) at 1:500 dilution + human spleen lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 127 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

CD11b antibody (N-term) (Cat. #AP10668a) immunohistochemistry analysis in formalin fixed and paraffin embedded human hepatocarcinoma followed by



peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the CD11b antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



CD11b Antibody (N-term) (Cat. #AP10668a) flow cytometric analysis of Jurkat cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

## Citations

- [Repetitive myocardial ischemia promotes coronary growth in the adult mammalian heart.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.