

# LMO4 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP10757C

## Product Information

---

<b>Application</b>	WB, IF, FC, E
<b>Primary Accession</b>	<a href="#">P61968</a>
<b>Other Accession</b>	<a href="#">P61969</a> , <a href="#">Q3SWZ8</a> , <a href="#">NP_006760.1</a>
<b>Reactivity</b>	Human, Mouse
<b>Predicted</b>	Bovine
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB24682
<b>Calculated MW</b>	17994
<b>Antigen Region</b>	104-131

## Additional Information

---

<b>Gene ID</b>	8543
<b>Other Names</b>	LIM domain transcription factor LMO4, Breast tumor autoantigen, LIM domain only protein 4, LMO-4, LMO4
<b>Target/Specificity</b>	This LMO4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 104-131 amino acids from the Central region of human LMO4.
<b>Dilution</b>	WB~~1:1000 IF~~1:10~50 FC~~1:10~50 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	LMO4 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	LMO4
<b>Function</b>	Transcription cofactor. Plays a role in establishing motor neuron identity, in

concert with MNX1, acting, at least in part, to disrupt LDB1-LHX3 complexes thereby negatively modulating interneuron genes in motor neurons.


## Background

This gene encodes a cysteine-rich protein that contains two LIM domains but lacks a DNA-binding homeodomain. The encoded protein may play a role as a transcriptional regulator or as an oncogene.


## References

Tian, Y., et al. Clin. Exp. Metastasis 27(7):455-463(2010)  
Ribases, M., et al. Biol. Psychiatry 66(10):926-934(2009)  
Montanez-Wiscovich, M.E., et al. Oncogene 28(41):3608-3618(2009)  
Huang, Z., et al. Dev. Biol. 327(1):132-142(2009)  
Caporaso, N., et al. PLoS ONE 4 (2), E4653 (2009) :


## Images



Fluorescent image of A549 cell stained with LMO4 Antibody (Center)(Cat#AP10757c).A549 cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then incubated with LMO4 primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400, 50 min at 37°C).Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7units/ml, 1 h at 37°C).LMO4 immunoreactivity is localized to Nucleus significantly.



LMO4 Antibody (Center) (Cat. #AP10757c) western blot analysis in mouse spleen tissue lysates (35ug/lane).This demonstrates the LMO4 antibody detected the LMO4 protein (arrow).



LMO4 Antibody (Center) (Cat. #AP10757c) flow cytometric analysis of MCF-7 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.