

# WDR41 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP10866B

## Product Information

---

Application	IHC-P, WB, E
Primary Accession	<a href="#">Q9HAD4</a>
Other Accession	<a href="#">NP_060738.2</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB28275
Calculated MW	51728
Antigen Region	249-277

## Additional Information

---

Gene ID	55255
Other Names	WD repeat-containing protein 41, WDR41
Target/Specificity	This WDR41 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 249-277 amino acids from the C-terminal region of human WDR41.
Dilution	IHC-P~~1:100 WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	WDR41 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	WDR41 ( <a href="#">HGNC:25601</a> )
Function	Non-catalytic component of the C9orf72-SMCR8 complex, a complex that has guanine nucleotide exchange factor (GEF) activity and regulates autophagy (PubMed: <a href="#">27103069</a> , PubMed: <a href="#">27193190</a> , PubMed: <a href="#">27617292</a> , PubMed: <a href="#">28195531</a> ). The C9orf72-SMCR8 complex promotes the exchange of


GDP to GTP, converting inactive GDP-bound RAB8A and RAB39B into their active GTP-bound form, thereby promoting autophagosome maturation (PubMed:[27103069](#)). As part of the C9orf72-SMCR8 complex, stimulates RAB8A and RAB11A GTPase activity in vitro, however WDR42 is shown not be an essential complex component for this function (PubMed:[32303654](#)). The C9orf72-SMCR8 complex also acts as a negative regulator of autophagy initiation by interacting with the ULK1/ATG1 kinase complex and inhibiting its protein kinase activity (PubMed:[27103069](#), PubMed:[27617292](#)).

**Cellular Location**                      Cytoplasm.


**References**

Bailey, S.D., et al. Diabetes Care (2010) In press :  
Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :  
Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)


**Images**



Immunohistochemical analysis of AP10866B on paraffin-embedded Human kidney tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Immunohistochemical analysis of AP10866B on paraffin-embedded Human placenta tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



All lanes : Anti-WDR41 Antibody (C-term) at 1:1000 dilution Lane 1: U-87 MG whole cell lysate Lane 2: U266B1 whole cell lysate Lane 3: Daudi whole cell lysate Lane 4: U-251 MG whole cell lysate Lane 5: 293 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 52 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

## Citations

---

- [Implementation of an antibody characterization procedure and application to the major ALS/FTD disease gene C9ORF72](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.