

BCL11B Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP10966c

Product Information

| | |
|-------------------|--|
| Application | WB, E |
| Primary Accession | Q9C0K0 |
| Other Accession | Q99PV8 , NP_612808.1 , NP_075049.1 |
| Reactivity | Human |
| Predicted | Mouse |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Clone Names | RB22239 |
| Calculated MW | 95519 |
| Antigen Region | 307-334 |

Additional Information

| | |
|--------------------|--|
| Gene ID | 64919 |
| Other Names | B-cell lymphoma/leukemia 11B, BCL-11B, B-cell CLL/lymphoma 11B, COUP-TF-interacting protein 2, Radiation-induced tumor suppressor gene 1 protein, hRit1, BCL11B, CTIP2, RIT1 |
| Target/Specificity | This BCL11B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 307-334 amino acids from the Central region of human BCL11B. |
| Dilution | WB~~1:1000 E~~Use at an assay dependent concentration. |
| Format | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Storage | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Precautions | BCL11B Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|----------|-------------|
| Name | BCL11B |
| Synonyms | CTIP2, RIT1 |

| | |
|--------------------------|--|
| Function | Key regulator of both differentiation and survival of T- lymphocytes during thymocyte development in mammals. Essential in controlling the responsiveness of hematopoietic stem cells to chemotactic signals by modulating the expression of the receptors CCR7 and CCR9, which direct the movement of progenitor cells from the bone marrow to the thymus (PubMed: 27959755). Is a regulator of IL2 promoter and enhances IL2 expression in activated CD4(+) T-lymphocytes (PubMed: 16809611). Tumor-suppressor that represses transcription through direct, TFCOUP2-independent binding to a GC-rich response element (By similarity). May also function in the P53-signaling pathway (By similarity). |
| Cellular Location | Nucleus. |
| Tissue Location | Highly expressed in brain and in malignant T-cell lines derived from patients with adult T-cell leukemia/lymphoma |


Background

This gene encodes a C2H2-type zinc finger protein and is closely related to BCL11A, a gene whose translocation may be associated with B-cell malignancies. The specific function of this gene has not yet been determined. Two alternatively spliced transcript variants, which encode distinct isoforms, have been reported.

References

Ganguli-Indra, G., et al. Exp. Dermatol. 18(11):994-996(2009)
Cherrier, T., et al. Oncogene 28(38):3380-3389(2009)
Ganguli-Indra, G., et al. PLoS ONE 4 (4), E5367 (2009) :
Cismasiu, V.B., et al. Virology 380(2):173-181(2008)
Desplats, P.A., et al. Neurobiol. Dis. 31(3):298-308(2008)

Images



All lanes : Anti-BCL11B Antibody (Center) at 1:16000 dilution
Lane 1: Jurkat whole cell lysate
Lane 2: MOLT-4 whole cell lysate
Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.
Predicted band size : 96 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.