

BCLAF1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP10988b

Product Information

Application WB, IHC-P, E Primary Accession Q9NYF8

Other Accession NP 055554.1, NP 001070908.1, NP 001070909.1

Reactivity Human
Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Clone Names RB25182
Calculated MW 106122
Antigen Region 658-687

Additional Information

Gene ID 9774

Other Names Bcl-2-associated transcription factor 1, Btf, BCLAF1, BTF, KIAA0164

Target/Specificity This BCLAF1 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 658-687 amino acids from the

C-terminal region of human BCLAF1.

Dilution WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions BCLAF1 Antibody (C-term) is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name BCLAF1

Synonyms BTF, KIAA0164

Function Death-promoting transcriptional repressor. May be involved in

cyclin-D1/CCND1 mRNA stability through the SNARP complex which

associates with both the 3'end of the CCND1 gene and its mRNA.

Cellular Location Cytoplasm. Nucleus. Nucleus speckle. Nucleus, nucleoplasm

Tissue Location Ubiquitous.


Background

This gene encodes a transcriptional repressor that interacts with several members of the BCL2 family of proteins. Overexpression of this protein induces apoptosis, which can be suppressed by co-expression of BCL2 proteins. The protein localizes to dot-like structures throughout the nucleus, and redistributes to a zone near the nuclear envelope in cells undergoing apoptosis. Multiple transcript variants encoding different isoforms have been found for this gene.


References

Sarras, H., et al. ScientificWorldJournal 10, 1450-1461 (2010): Renert, A.F., et al. J. Proteome Res. 8(10):4810-4822(2009) Liu, H., et al. Mol. Cell. Biol. 27(24):8480-8491(2007) Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007) Olsen, J.V., et al. Cell 127(3):635-648(2006)

Images



BCLAF1 Antibody (C-term) (Cat. #AP10988b) western blot analysis in MCF-7 cell line lysates (35ug/lane). This demonstrates the BCLAF1 antibody detected the BCLAF1 protein (arrow).



BCLAF1 antibody (C-term) (Cat. #AP10988b) immunohistochemistry analysis in formalin fixed and paraffin embedded human skeletal muscle followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the BCLAF1 antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.