10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

HDAC11 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1111b

Product Information

Application	WB, IHC-P, E
Primary Accession	Q96DB2
Reactivity	Human, Mouse
Predicted	Bovine, Canine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	39183
Antigen Region	313-345

Additional Information

Gene ID	79885
Other Names	Histone deacetylase 11, HD11, HDAC11
Target/Specificity	This HDAC11 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 313-345 amino acids from the C-terminal region of human HDAC11.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	HDAC11 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	HDAC11
Function	Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes.

Cellular Location	Nucleus.
Tissue Location	Weakly expressed in most tissues. Strongly expressed in brain, heart, skeletal muscle, kidney and testis


Background

HDAC11 is responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. The predominantly nuclear HDAC11, which interacts with HDAC6, is weakly expressed in most tissues, and strongly expressed in brain, heart, skeletal muscle, kidney and testis. Its activity is inhibited by trapoxin, a known histone deacetylase inhibitor.


References

Keedy, K.S. et al. J Virol. May; 83(10): 4749?756(2009).
Voelter-Mahlknecht S, et al., Int J Mol Med. 2005 Oct;16(4):589-98.
Bradbury CA, et al., Leukemia. 2005 Oct;19(10):1751-9.
Gregoretti IV, et al., J Mol Biol. 2004 Apr 16;338(1):17-31.
Gao, L., et al., J. Biol. Chem. 277(28):25748-25755 (2002).

Images



The anti-HDAC11 Pab (Cat. #AP1111b) is used in Western blot to detect HDAC11 in mouse brain tissue lysate.



Formalin-fixed and paraffin-embedded human testicle tumor tissue reacted with HDAC11 antibody (C-term)(Cat.#AP1111b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Citations

- The functional interactome landscape of the human histone deacetylase family.
- A limited group of class I histone deacetylases acts to repress human immunodeficiency virus type 1 expression.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.