

UNCX Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP11164b

Product Information

Application	WB, IF, E
Primary Accession	A6NJT0
Other Accession	NP_001073930.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB28877
Calculated MW	53690
Antigen Region	382-410

Additional Information

Gene ID	340260
Other Names	Homeobox protein unc-4 homolog, Homeobox protein Uncx41, UNCX, UNCX41
Target/Specificity	This UNCX antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 382-410 amino acids from the C-terminal region of human UNCX.
Dilution	WB~~1:1000 IF~~1:25 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	UNCX Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	UNCX
Synonyms	UNCX4.1
Function	Transcription factor involved in somitogenesis and neurogenesis. Required

for the maintenance and differentiation of particular elements of the axial skeleton. May act upstream of PAX9. Plays a role in controlling the development of connections of hypothalamic neurons to pituitary elements, allowing central neurons to reach the peripheral blood circulation and to deliver hormones for control of peripheral functions (By similarity).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108}.


Background

Myosins are actin-based motor molecules with ATPase activity. Unconventional myosins serve in intracellular movements. Their highly divergent tails are presumed to bind to membranous compartments, which would be moved relative to actin filaments (By similarity).


References

Venter, J.C., et al. Science 291(5507):1304-1351(2001)

Images



Immunofluorescent analysis of U251 cells, using UNCX Antibody (C-term) (Cat. #AP11164b). AP11164b was diluted at 1:25 dilution. Alexa Fluor 488-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody (green). Cytoplasmic actin was counterstained with Dylight Fluor® 554 (red) conjugated Phalloidin (red).



UNCX Antibody (C-term) (Cat. #AP11164b) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the UNCX antibody detected the UNCX protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.