

GOG8A Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP11196A

Product Information

Application	WB, IHC-P, IF, E
Primary Accession	A7E2F4
Other Accession	A8MQT2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB30142
Calculated MW	70117
Antigen Region	129-157

Additional Information


Gene ID	23015
Other Names	Golgin subfamily A member 8A, 88 kDa Golgi matrix protein, GM88, GM88 autoantigen, GOLGA8A
Target/Specificity	This GOG8A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 129-157 amino acids from the N-terminal region of human GOG8A.
Dilution	WB~~1:1000 IHC-P~~1:100~500 IF~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	GOG8A Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	GOLGA8A
Synonyms	KIAA0855

Function	May be involved in maintaining Golgi structure.
Cellular Location	Golgi apparatus, Golgi stack membrane; Peripheral membrane protein


Images



GOG8A Antibody (N-term) (Cat. #AP11196a) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the GOG8A antibody detected the GOG8A protein (arrow).



GOG8A Antibody (N-term) (Cat. #AP11196a) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of GOG8A Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Confocal immunofluorescent analysis of GOG8A Antibody (N-term) (Cat#AP11196a) with 293 cells followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.