

GPX3 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP11221c

Product Information

Application	WB, IHC-P, E
Primary Accession	P22352
Other Accession	NP_002075
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB18898
Calculated MW	25552
Antigen Region	93-123

Additional Information

Gene ID	2878
Other Names	Glutathione peroxidase 3, GPx-3, GSHPx-3, Extracellular glutathione peroxidase, Plasma glutathione peroxidase, GPx-P, GSHPx-P, GPX3, GPXP
Target/Specificity	This GPX3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 93-123 amino acids from the Central region of human GPX3.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	GPX3 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	GPX3 (HGNC:4555)
Synonyms	GPXP
Function	Protects cells and enzymes from oxidative damage, by catalyzing the

reduction of hydrogen peroxide, lipid peroxides and organic hydroperoxide, by glutathione.

Cellular Location Secreted.

Tissue Location Secreted in plasma.


Background

This gene product belongs to the glutathione peroxidase family, which functions in the detoxification of hydrogen peroxide. It contains a selenocysteine (Sec) residue at its active site. The selenocysteine is encoded by the UGA codon, which normally signals translation termination. The 3' UTR of Sec-containing genes have a common stem-loop structure, the sec insertion sequence (SECIS), which is necessary for the recognition of UGA as a Sec codon rather than as a stop signal.

References

Wang, J.Y., et al. Clin. Chim. Acta 411 (19-20), 1432-1436 (2010) : Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010) Fullerton, J.M., et al. Bipolar Disord 12(5):550-556(2010) Wang, Y., et al. J. Hum. Genet. 55(8):490-494(2010) Mistry, H.D., et al. Placenta 31(5):401-408(2010)

Images



All lanes : Anti-GPX3 Antibody (Center) at 1:1000 dilution
Lane 1: M. plasma lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Observed band size : 22kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Citations

- [Silencing GPX3 Expression Promotes Tumor Metastasis in Human Thyroid Cancer.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.