

GGH Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP11236a

Product Information

Application	IHC-P, FC, WB, E
Primary Accession	Q92820
Other Accession	NP_003869
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB20066
Calculated MW	35964
Antigen Region	7-34

Additional Information

Gene ID	8836
Other Names	Gamma-glutamyl hydrolase, Conjugase, GH, Gamma-Glu-X carboxypeptidase, GGH
Target/Specificity	This GGH antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 7-34 amino acids from the N-terminal region of human GGH.
Dilution	IHC-P~~1:100~500 FC~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	GGH Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	GGH (HGNC:4248)
Function	Hydrolyzes the polyglutamate sidechains of pteroylpolyglutamates. Progressively removes gamma-glutamyl residues from

pteroylpoly-gamma-glutamate to yield pteroyl-alpha-glutamate (folic acid) and free glutamate (PubMed:[11005824](#), PubMed:[8816764](#)). May play an important role in the bioavailability of dietary pteroylpolyglutamates and in the metabolism of pteroylpolyglutamates and antifolates.

Cellular Location

Secreted, extracellular space. Lysosome. Melanosome. Note=While its intracellular location is primarily the lysosome, most of the enzyme activity is secreted Identified by mass spectrometry in melanosome fractions from stage I to stage IV.


Background

This gene catalyzes the hydrolysis of folylpoly-gamma-glutamates and antifolylpoly-gamma-glutamates by the removal of gamma-linked polyglutamates and glutamate. [provided by RefSeq].


References

Adjei, A.A., et al. J Thorac Oncol 5(9):1346-1353(2010)
Liu, C.Y., et al. Carcinogenesis 31(7):1259-1263(2010)
Organista-Nava, J., et al. Leuk. Res. 34(6):728-732(2010)
Figueiredo, J.C., et al. Cancer Causes Control 21(4):597-608(2010)
Dervieux, T., et al. Pharmacogenet. Genomics (2009) In press :

Images




Anti-GGH Antibody (N-term) at 1:2000 dilution + U-87 MG whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 36 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



GGH Antibody (N-term) (Cat. #AP11236a) immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of GGH Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

GGH Antibody (N-term) (Cat. #AP11236a) flow cytometric analysis of HL-60 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.