

FA96A Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP11250b

Product Information

Application	WB, IHC-P, IF, E
Primary Accession	Q9H5X1
Other Accession	Q9DCL2 , Q3T0U7 , NP_115607.1
Reactivity	Human, Rat, Mouse
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB28646
Calculated MW	18355
Antigen Region	109-138

Additional Information

Gene ID	84191
Other Names	MIP18 family protein FAM96A, FAM96A
Target/Specificity	This FA96A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 109-138 amino acids from the C-terminal region of human FA96A.
Dilution	WB~~1:1000 IHC-P~~1:100~500 IF~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	FA96A Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CIAO2A (HGNC:26235)
Function	Component of the cytosolic iron-sulfur protein assembly (CIA) complex, a multiprotein complex that mediates the incorporation of iron-sulfur cluster

into extramitochondrial Fe/S proteins (PubMed:[23891004](#)). As a CIA complex component and in collaboration with CIAO1 specifically matures ACO1 and stabilizes IREB2, connecting cytosolic iron-sulfur protein maturation with cellular iron regulation (PubMed:[23891004](#)). May play a role in chromosome segregation through establishment of sister chromatid cohesion. May induce apoptosis in collaboration with APAF1 (PubMed:[25716227](#)).


Cellular Location Cytoplasm.

Tissue Location Substantially enriched in macrophages.

References

Gerhard, D.S., et al. Genome Res. 14 (10B), 2121-2127 (2004) :

Images



Anti-FA96A Antibody (C-term) at 1:2000 dilution + K562 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 18 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.