

PSMB11 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP11339b

Product Information

Application	WB, IHC-P, E
Primary Accession	A5LHX3
Other Accession	NP_001093250.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB28790
Calculated MW	32530
Antigen Region	250-278

Additional Information

Gene ID	122706
Other Names	Proteasome subunit beta type-11, Proteasome subunit beta-5t, PSMB11
Target/Specificity	This PSMB11 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 250-278 amino acids from the C-terminal region of human PSMB11.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PSMB11 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PSMB11
Function	The proteasome is a multicatalytic proteinase complex which is characterized by its ability to cleave peptides with Arg, Phe, Tyr, Leu, and Glu adjacent to the leaving group at neutral or slightly basic pH. The proteasome has an ATP-dependent proteolytic activity. Incorporated instead of PSMB5 or

PSMB8, this unit reduces the chymotrypsin-like activity of the proteasome (By similarity). Plays a pivotal role in development of CD8-positive T cells (By similarity).

Cellular Location

Cytoplasm {ECO:0000255 | PROSITE-ProRule:PRU00809}. Nucleus


Background

Proteasomes generate peptides that are presented by major histocompatibility complex (MHC) I molecules to other cells of the immune system. Proteolysis is conducted by 20S proteasomes, complexes of 28 subunits arranged as a cylinder in 4 heteroheptameric rings: alpha-1 to -7, beta-1 to -7, beta-1 to -7, and alpha-1 to -7. The catalytic subunits are beta-1 (PSMB6; MIM 600307), beta-2 (PSMB7; MIM 604030), and beta-5 (PSMB5; MIM 600306). Three additional subunits, beta-1i (PSMB9; MIM 177045), beta-2i (PSMB10; MIM 176847), and beta-5i (PSMB8; MIM 177046), are induced by gamma-interferon (IFNG; MIM 147570) and are preferentially incorporated into proteasomes to make immunoproteasomes. PSMB11, or beta-5t, is a catalytic subunit expressed exclusively in cortical thymic epithelial cells (Murata et al., 2007 [PubMed 17540904]).


References

Tomaru, U., et al. Blood 113(21):5186-5191(2009)
Murata, S., et al. Science 316(5829):1349-1353(2007)

Images



PSMB11 Antibody (C-term) (Cat. #AP11339b) immunohistochemistry analysis in formalin fixed and paraffin embedded human thymoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of PSMB11 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



All lanes : Anti-PSMB11 Antibody (C-term) at 1:1000 dilution+ 3T3-L1 cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size : 38kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.