10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

C9JLR9 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11349b

Product Information

Application	WB, IHC-P, IF, E
Primary Accession	C9JLR9
Other Accession	Q4VA45, NP_001138408.1
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB29021
Calculated MW	73197
Antigen Region	556-584

Additional Information

Gene ID	65998
Other Names	Uncharacterized protein C11orf95, C11orf95
Target/Specificity	This C9JLR9 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 556-584 amino acids from the C-terminal region of human C9JLR9.
Dilution	WB~~1:1000 IHC-P~~1:100~500 IF~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	C9JLR9 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Name	ZFTA (HGNC:28449)
Synonyms	C11orf95


References

Ota, T., et al. Nat. Genet. 36(1):40-45(2004)


Images



All lanes: Anti-C9JLR9 Antibody (C-term) at 1:2000 dilution Lane 1: HepG2 whole cell lysate Lane 2: PC-3 whole cell lysate Lane 3: HT-29 whole cell lysate Lane 4: Rat ovary lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 73 KDa Blocking/Dilution buffer: 5% NFDM/TBST.



C9JLR9 Antibody (C-term) (Cat. #AP11349b)immunohistochemistry analysis in formalin fixed and paraffin embedded human adipose tissue followed by peroxidase conjugation of the secondary antibody and DAB staining.This data demonstrates the use of C9JLR9 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Confocal immunofluorescent analysis of C9JLR9 Antibody (C-term)(Cat#AP11349b) with HepG2 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit lgG (green). Actin filaments have been labeled with Alexa Fluor555 phalloidin (red). DAPI was used to stain the cell nuclear (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.