

WDR70 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP11355B

Product Information

Application	WB, IHC-P, FC, IF, E
Primary Accession	Q9NW82
Other Accession	Q5EB92 , Q3TWF6 , Q32LB0 , NP_060504.1
Reactivity	Human, Rat, Mouse
Predicted	Mouse, Rat, Bovine, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB29043
Calculated MW	73201
Antigen Region	575-604

Additional Information

Gene ID	55100
Other Names	WD repeat-containing protein 70, WDR70
Target/Specificity	This WDR70 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 575-604 amino acids from the C-terminal region of human WDR70.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 IF~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	WDR70 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Name	WDR70
-------------	-------


References

Olsen, J.V., et al. Cell 127(3):635-648(2006)
Ota, T., et al. Nat. Genet. 36(1):40-45(2004)


Images




Anti-WDR70 Antibody (C-term) at 1:500 dilution + HL-60 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 70 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



WDR70 Antibody (C-term) (Cat. #AP11355b) immunohistochemistry analysis in formalin fixed and paraffin embedded human esophageal carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of WDR70 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



WDR70 Antibody (C-term) (Cat. #AP11355b) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Confocal immunofluorescent analysis of WDR70 Antibody (C-term) (Cat. #AP11355b) with Hela cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.