

# STRA8 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP11436b

## Product Information

---

<b>Application</b>	IHC-P-Leica, WB, E
<b>Primary Accession</b>	<a href="#">Q7Z7C7</a>
<b>Other Accession</b>	<a href="#">NP_872295.1</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB29578
<b>Calculated MW</b>	36908
<b>Antigen Region</b>	286-315

## Additional Information

---

<b>Gene ID</b>	346673
<b>Other Names</b>	Stimulated by retinoic acid gene 8 protein homolog, STRA8 {ECO:0000312   EMBL:AAP471631}
<b>Target/Specificity</b>	This STRA8 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 286-315 amino acids from the C-terminal region of human STRA8.
<b>Dilution</b>	IHC-P-Leica~~1:500 WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	STRA8 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	STRA8 ( <a href="#">HGNC:30653</a> )
<b>Function</b>	Meiosis-inducer required for the transition into meiosis for both female and male germ cells. In female germ cells, acts downstream of ZGLP1 as a key

effector of the meiotic program: required for premeiotic DNA replication and subsequent events in meiotic prophase. During spermatogenesis, next to its role in meiotic initiation, promotes (but is not required for) spermatogonial differentiation. In complex with MEIOSIN, directly activates the transcription of a subset of critical meiotic genes playing a central role in cell-cycle switching from mitosis to meiosis.

#### Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:P70278}. Nucleus {ECO:0000250|UniProtKB:P70278}. Note=Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent {ECO:0000250|UniProtKB:P70278}

#### Tissue Location

Expressed specifically in testis and fetal ovaries.


## Background

This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in humans.


## References

- Le Bouffant, R., et al. Hum. Reprod. 25(10):2579-2590(2010)  
Aston, K.I., et al. Hum. Reprod. 25(6):1383-1397(2010)  
Chen, B., et al. Asian J. Androl. 11(5):557-565(2009)  
Houmard, B., et al. Biol. Reprod. 81(2):438-443(2009)  
Stouffs, K., et al. Verh. K. Acad. Geneesk. Belg. 71(3):115-139(2009)

## Images



Immunohistochemical analysis of AP11436B on paraffin-embedded human lung tissue was performed on the Leica® BOND RXm. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:250) for 15min at room temperature. Leica Bond Polymer Refine Detection was used as the secondary antibody.



All lanes : Anti-STRA8 Antibody (C-term) at 1:2000 dilution  
Lane 1: 293 whole cell lysate Lane 2: LNCaP whole cell lysate Lane 3: PC-3 whole cell lysate Lane 4: DU145 whole cell lysate Lane 5: SK-OV-3 whole cell lysate Lane 6: OVCAR3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 37 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.