

MESDC2 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP11638b

Product Information

Application	IHC-P, IF, WB, E
Primary Accession	Q14696
Other Accession	NP_055969.1
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB29835
Calculated MW	26077
Antigen Region	206-234

Additional Information

Gene ID	23184
Other Names	LDLR chaperone MESD, Mesoderm development candidate 2, Mesoderm development protein, Renal carcinoma antigen NY-REN-61, MESDC2, KIAA0081, MESD
Target/Specificity	This MESDC2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 206-234 amino acids from the C-terminal region of human MESDC2.
Dilution	IHC-P~~1:100~500 IF~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	MESDC2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	MESD (HGNC:13520)
Synonyms	KIAA0081, MESDC2, MESDM

Function	Chaperone specifically assisting the folding of beta- propeller/EGF modules within the family of low-density lipoprotein receptors (LDLRs) (PubMed: 15014448). Acts as a modulator of the Wnt pathway through chaperoning the coreceptors of the canonical Wnt pathway, LRP5 and LRP6, to the plasma membrane (PubMed: 17488095 , PubMed: 23572575). Essential for specification of embryonic polarity and mesoderm induction. Plays an essential role in neuromuscular junction (NMJ) formation by promoting cell-surface expression of LRP4 (By similarity). May regulate phagocytosis of apoptotic retinal pigment epithelium (RPE) cells (By similarity).
Cellular Location	Endoplasmic reticulum Note=Released from apoptotic cells and shed photoreceptor outer segments. {ECO:0000250 UniProtKB:Q9ERE7}


Background

Chaperone specifically assisting the folding of beta-propeller/EGF modules within the family of low-density lipoprotein receptors (LDLRs). Acts as a modulator of the Wnt pathway through chaperoning the coreceptors of the canonical Wnt pathway, LRP5 and LRP6, to the plasma membrane. Essential for specification of embryonic polarity and mesoderm induction.


References

Murrills, R.J., et al. J. Cell. Biochem. 108(5):1066-1075(2009)
 Li, Y., et al. FEBS Lett. 580(22):5423-5428(2006)
 Veltman, I.M., et al. Hum. Mol. Genet. 14(14):1955-1963(2005)
 Clark, H.F., et al. Genome Res. 13(10):2265-2270(2003)
 Hsieh, J.C., et al. Cell 112(3):355-367(2003)


Images



Western blot analysis of lysates from K562, mouse NIH/3T3 cell line (from left to right), using MESDC2 Antibody (C-term) (Cat. #AP11638b). AP11638b was diluted at 1:1000 at each lane. A goat anti-rabbit(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.



MESDC2 Antibody (C-term)(Cat. #AP11638b)immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue followed by peroxidase conjugation of the secondary antibody and DAB staining.This data demonstrates the use of MESDC2 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



Confocal immunofluorescent analysis of MESDC2 Antibody (C-term)(Cat#AP11638b) with 293 cell followed by Alexa Fluor[®] 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.