

## KHSRP Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP11709c

### Product Information

---

<b>Application</b>	IHC-P, IF, WB, E
<b>Primary Accession</b>	<a href="#">Q92945</a>
<b>Other Accession</b>	<a href="#">Q99PF5</a> , <a href="#">Q3U0V1</a> , <a href="#">NP_003676.2</a>
<b>Reactivity</b>	Human
<b>Predicted</b>	Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB20821
<b>Calculated MW</b>	73115
<b>Antigen Region</b>	470-498

### Additional Information

---

<b>Gene ID</b>	8570
<b>Other Names</b>	Far upstream element-binding protein 2, FUSE-binding protein 2, KH type-splicing regulatory protein, KSRP, p75, KHSRP, FUBP2
<b>Target/Specificity</b>	This KHSRP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 470-498 amino acids from the Central region of human KHSRP.
<b>Dilution</b>	IHC-P~~1:100~500 IF~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	KHSRP Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

### Protein Information

---

<b>Name</b>	KHSRP
<b>Synonyms</b>	FUBP2

<b>Function</b>	Binds to the dendritic targeting element and may play a role in mRNA trafficking (By similarity). Part of a ternary complex that binds to the downstream control sequence (DCS) of the pre-mRNA. Mediates exon inclusion in transcripts that are subject to tissue- specific alternative splicing. May interact with single-stranded DNA from the far-upstream element (FUSE). May activate gene expression. Also involved in degradation of inherently unstable mRNAs that contain AU-rich elements (AREs) in their 3'-UTR, possibly by recruiting degradation machinery to ARE-containing mRNAs.
<b>Cellular Location</b>	Nucleus. Cytoplasm. Note=A small proportion is also found in the cytoplasm of neuronal cell bodies and dendrites.
<b>Tissue Location</b>	Detected in neural and non-neural cell lines.


## Background

The KHSRP gene encodes a multifunctional RNA-binding protein implicated in a variety of cellular processes, including transcription, alternative pre-mRNA splicing, and mRNA localization (Min et al., 1997 [PubMed 9136930]; Gherzi et al., 2004 [PubMed 15175153]).


## References

Michlewski, G., et al. Nat. Struct. Mol. Biol. 17(8):1011-1018(2010)  
 Bikkavilli, R.K., et al. J. Cell. Sci. 123 (PT 8), 1352-1362 (2010) :  
 Malz, M., et al. Hepatology 50(4):1130-1139(2009)  
 Trabucchi, M., et al. Nature 459(7249):1010-1014(2009)  
 Nechama, M., et al. BMC Cell Biol. 10, 70 (2009) :


## Images



Anti-KHSRP Antibody (Center) at 1:1000 dilution + HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 73 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



KHSRP Antibody (Center) (Cat. #AP11709c) immunohistochemistry analysis in formalin fixed and paraffin embedded human testis tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of KHSRP Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



Confocal immunofluorescent analysis of KHSRP Antibody (Center)(Cat#AP11709c) with U-251MG cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 phalloidin (red). DAPI was used to stain the cell nuclear (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.