

# S100A1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP11779b

## Product Information

---

<b>Application</b>	WB, IHC-P, E
<b>Primary Accession</b>	<a href="#">P23297</a>
<b>Other Accession</b>	<a href="#">P02639</a> , <a href="#">NP_006262.1</a>
<b>Reactivity</b>	Human
<b>Predicted</b>	Bovine
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB19108
<b>Calculated MW</b>	10546
<b>Antigen Region</b>	45-75

## Additional Information

---

<b>Gene ID</b>	6271
<b>Other Names</b>	Protein S100-A1, S-100 protein alpha chain, S-100 protein subunit alpha, S100 calcium-binding protein A1, S100A1, S100A
<b>Target/Specificity</b>	This S100A1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 45-75 amino acids from the C-terminal region of human S100A1.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	S100A1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	S100A1
<b>Synonyms</b>	S100A

<b>Function</b>	Small calcium binding protein that plays important roles in several biological processes such as Ca(2+) homeostasis, chondrocyte biology and cardiomyocyte regulation (PubMed: <a href="#">12804600</a> ). In response to an increase in intracellular Ca(2+) levels, binds calcium which triggers conformational changes (PubMed: <a href="#">23351007</a> ). These changes allow interactions with specific target proteins and modulate their activity (PubMed: <a href="#">22399290</a> ). Regulates a network in cardiomyocytes controlling sarcoplasmic reticulum Ca(2+) cycling and mitochondrial function through interaction with the ryanodine receptors RYR1 and RYR2, sarcoplasmic reticulum Ca(2+)-ATPase/ATP2A2 and mitochondrial F1-ATPase (PubMed: <a href="#">12804600</a> ). Facilitates diastolic Ca(2+) dissociation and myofilament mechanics in order to improve relaxation during diastole (PubMed: <a href="#">11717446</a> ).
<b>Cellular Location</b>	Cytoplasm. Sarcoplasmic reticulum. Mitochondrion {ECO:0000250 UniProtKB:P56565}
<b>Tissue Location</b>	Highly prevalent in heart (PubMed:12804600, PubMed:1384693). Also found in lesser quantities in skeletal muscle and brain (PubMed:1384693).


## Background

The protein encoded by this gene is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, and involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21. This protein may function in stimulation of Ca<sup>2+</sup>-induced Ca<sup>2+</sup> release, inhibition of microtubule assembly, and inhibition of protein kinase C-mediated phosphorylation. Reduced expression of this protein has been implicated in cardiomyopathies.

## References


Marlatt, N.M., et al. Protein Expr. Purif. 73(1):58-64(2010)  
van Dieck, J., et al. FEBS Lett. 584(15):3269-3274(2010)  
Sviatoha, V., et al. Melanoma Res. 20(2):118-125(2010)  
DeRycke, M.S., et al. Am. J. Clin. Pathol. 132(6):846-856(2009)  
Yusenko, M.V., et al. Int. J. Biol. Sci. 5(6):517-527(2009)

## Images



S100A1 Antibody (C-term) (Cat. #AP11779b) western blot analysis in MDA-MB435 cell line lysates (35ug/lane). This demonstrates the S100A1 antibody detected the S100A1 protein (arrow).

S100A1 Antibody (C-term) (Cat. #AP11779b) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary



antibody and DAB staining. This data demonstrates the use of S100A1 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.