

TNFAIP8L3 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP11822c

Product Information

Application WB, E **Primary Accession Q5GJ75** Other Accession NP 997264.2 Reactivity Human Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB29747 Calculated MW 32659 136-164 **Antigen Region**

Additional Information

Gene ID 388121

Other Names Tumor necrosis factor alpha-induced protein 8-like protein 3, TNF

alpha-induced protein 8-like protein 3, TNFAIP8-like protein 3, TNFAIP8L3

Target/SpecificityThis TNFAIP8L3 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 136-164 amino acids from the Central

region of human TNFAIP8L3.

Dilution WB~~1:1000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions TNFAIP8L3 Antibody (Center) is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name TNFAIP8L3

Synonyms TIPE3

Function Acts as a lipid transfer protein. Preferentially captures and shuttles two lipid

second messengers, i.e., phosphatidylinositol 4,5- bisphosphate and phosphatidylinositol 3,4,5-trisphosphate and increases their levels in the plasma membrane. Additionally, may also function as a lipid-presenting protein to enhance the activity of the PI3K-AKT and MEK-ERK pathways. May act as a regulator of tumorigenesis through its activation of phospholipid signaling.

Cellular Location Cytoplasm. Cell membrane {ECO:0000250 | UniProtKB:Q3TBL6}. Note=On

PDGF activation, translocates from cytoplasm to plasma membrane

{ECO:0000250 | UniProtKB:Q3TBL6}


Tissue Location Widely expressed (at protein level) (PubMed:25479791). Highly expressed in

most carcinoma cell lines (PubMed:25242044, PubMed:25479791).

References

Strausberg, R.L., et al. Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903(2002)

Images



TNFAIP8L3 Antibody (Center) (Cat. #AP11822c) western blot analysis in MDA-MB231 cell line lysates (35ug/lane). This demonstrates the TNFAIP8L3 antibody detected the TNFAIP8L3 protein (arrow).

Citations

• Expression of TIPE family members in human colorectal cancer

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.