10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

WTIP Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11837b

Product Information

Application	WB, IHC-P, IF, E
Primary Accession	A6NIX2
Other Accession	Q7TQJ8
Reactivity	Human, Mouse
Predicted	Mouse, Rat, Bovine, Canine, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB29854
Calculated MW	45124
Antigen Region	243-271

Additional Information

Gene ID	126374
Other Names	Wilms tumor protein 1-interacting protein, WT1-interacting protein, WTIP
Target/Specificity	This WTIP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 243-271 amino acids of human WTIP.
Dilution	WB~~1:500 IHC-P~~1:100~500 IF~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	WTIP Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	WTIP
Function	Adapter or scaffold protein which participates in the assembly of numerous protein complexes and is involved in several cellular processes such as cell fate determination, cytoskeletal organization, repression of gene

transcription, cell-cell adhesion, cell differentiation, proliferation and migration. Positively regulates microRNA (miRNA)-mediated gene silencing. Negatively regulates Hippo signaling pathway and antagonizes phosphorylation of YAP1. Acts as a transcriptional corepressor for SNAI1 and SNAI2/SLUG-dependent repression of E-cadherin transcription. Acts as a hypoxic regulator by bridging an association between the prolyl hydroxylases and VHL enabling efficient degradation of HIF1A. In podocytes, may play a role in the regulation of actin dynamics and/or foot process cytoarchitecture (By similarity). In the course of podocyte injury, shuttles into the nucleus and acts as a transcription regulator that represses WT1-dependent transcription regulation, thereby translating changes in slit diaphragm structure into altered gene expression and a less differentiated phenotype. Involved in the organization of the basal body (By similarity). Involved in cilia growth and positioning (By similarity).


Cellular Location

Cell junction, adherens junction. Nucleus. Cytoplasm, P-body. Note=Following podocyte injury, caused by treatment with LPS, puromycin aminonucleoside, ultraviolet or hydrogen peroxide, translocates from sites of cell-cell contacts into the cytosol and nucleus. The shift from cell contacts to intracellular plaques starts as early as 1 hour after LPS stimulation and intranuclear localization begins 3 hours after LPS treatment. Maximal nuclear localization is achieved 6 hours after LPS treatment. Nuclear translocation requires dynein motor activity and intact microtubule network (By similarity). Returns to cell-cell contacts 24 hours after LPS stimulation. In the presence of ROR2, localizes to the plasma membrane (By similarity).

Background


WTIP may monitor slit diaphragm protein assembly, a specialized adherens junction characteristic of podocytes. In case of podocyte injury, it shuttles into the nucleus and acts as a transcription regulator that represses WT1-dependent transcription regulation, thereby translating changes in slit diaphragm structure into altered gene expression and a less differentiated phenotype (By similarity).


Images



WTIP Antibody (C-term) (Cat. #AP11837b)immunohistochemistry analysis in formalin fixed and paraffin embedded human hepatocarcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining.This data demonstrates the use of WTIP Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

WTIP Antibody (C-term) (Cat. #AP11837b) western blot analysis in NCI-H460,WiDr cell line lysates (35ug/lane).This demonstrates the WTIP antibody detected the WTIP protein (arrow).





Confocal immunofluorescent analysis of WTIP Antibody (C-term)(Cat#AP11837b) with WiDr cell followed by Alexa Fluor 488-conjugated goat anti-rabbit lgG (green). DAPI was used to stain the cell nuclear (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.