

TRADD Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP11963c

Product Information

Application	WB, IHC-P, FC, IF, E
Primary Accession	Q15628
Other Accession	NP_003780
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB18788
Calculated MW	34247
Antigen Region	126-151

Additional Information

Gene ID	8717
Other Names	Tumor necrosis factor receptor type 1-associated DEATH domain protein, TNFR1-associated DEATH domain protein, TNFRSF1A-associated via death domain, TRADD
Target/Specificity	This TRADD antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 126-151 amino acids from the Central region of human TRADD.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 IF~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TRADD Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TRADD {ECO:0000303 PubMed:7758105, ECO:0000312 HGNC:HGNC:12030}
Function	Adapter molecule for TNFRSF1A/TNFR1 that specifically associates with the

cytoplasmic domain of activated TNFRSF1A/TNFR1 mediating its interaction with FADD (PubMed:[23955153](#), PubMed:[7758105](#), PubMed:[8612133](#)). Overexpression of TRADD leads to two major TNF-induced responses, apoptosis and activation of NF-kappa-B (PubMed:[7758105](#), PubMed:[8612133](#)). The nuclear form acts as a tumor suppressor by preventing ubiquitination and degradation of isoform p19ARF/ARF of CDKN2A by TRIP12: acts by interacting with TRIP12, leading to disrupt interaction between TRIP12 and isoform p19ARF/ARF of CDKN2A (By similarity).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q3U0V2}. Cytoplasm. Cytoplasm, cytoskeleton. Note=Shuttles between the cytoplasm and the nucleus. {ECO:0000250|UniProtKB:Q3U0V2}

Tissue Location

Found in all examined tissues.


Background

The protein encoded by this gene is a death domain containing adaptor molecule that interacts with TNFRSF1A/TNFR1 and mediates programmed cell death signaling and NF-kappaB activation. This protein binds adaptor protein TRAF2, reduces the recruitment of inhibitor-of-apoptosis proteins (IAPs) by TRAF2, and thus suppresses TRAF2 mediated apoptosis. This protein can also interact with receptor TNFRSF6/FAS and adaptor protein FADD/MORT1, and is involved in the Fas-induced cell death pathway. [provided by RefSeq].

References



Pointon, J.J., et al. Ann. Rheum. Dis. 69(6):1243-1246(2010)
Hosgood, H.D. III, et al. Occup Environ Med 66(12):848-853(2009)
Yerges, L.M., et al. J. Bone Miner. Res. 24(12):2039-2049(2009)
Liang, X.S., et al. Br. J. Haematol. 146(4):418-423(2009)
Dagle, J.M., et al. Pediatrics 123(4):1116-1123(2009)

Images




TRADD Antibody (Center) (Cat. #AP11963c) western blot analysis in HeLa cell line lysates (35ug/lane). This demonstrates the TRADD antibody detected the TRADD protein (arrow).

TRADD Antibody (Center) (Cat. #AP11963c) immunohistochemistry analysis in formalin fixed and paraffin embedded human breast carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of TRADD Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



TRADD Antibody (Center) (Cat. #AP11963c) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.



Confocal immunofluorescent analysis of TRADD Antibody (Center) (Cat#AP11963c) with Hela cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). Actin filaments have been labeled with Alexa Fluor 555 phalloidin (red). DAPI was used to stain the cell nuclear (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.