

# MSI1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP11965a

## Product Information

---

<b>Application</b>	IHC-P, FC, WB, E
<b>Primary Accession</b>	<a href="#">Q43347</a>
<b>Other Accession</b>	<a href="#">Q8K3P4</a> , <a href="#">Q61474</a> , <a href="#">NP_002433</a>
<b>Reactivity</b>	Human
<b>Predicted</b>	Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB19177
<b>Calculated MW</b>	39125
<b>Antigen Region</b>	67-93

## Additional Information

---

<b>Gene ID</b>	4440
<b>Other Names</b>	RNA-binding protein Musashi homolog 1, Musashi-1, MSI1
<b>Target/Specificity</b>	This MSI1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 67-93 amino acids from the N-terminal region of human MSI1.
<b>Dilution</b>	IHC-P~~1:100~500 FC~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	MSI1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	MSI1
<b>Function</b>	RNA binding protein that regulates the expression of target mRNAs at the translation level. Regulates expression of the NOTCH1 antagonist NUMB.

Binds RNA containing the sequence 5'-GUUAGUUAGUUAGUU- 3' and other sequences containing the pattern 5'-[GA]U(1-3)AGU-3'. May play a role in the proliferation and maintenance of stem cells in the central nervous system (By similarity).

#### Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q61474}. Nucleus {ECO:0000250|UniProtKB:Q61474}

#### Tissue Location

Detected in fetal kidney, brain, liver and lung, and in adult brain and pancreas. Detected in hepatoma cell lines


## Background

This gene encodes a protein containing two conserved tandem RNA recognition motifs. Similar proteins in other species function as RNA-binding proteins and play central roles in posttranscriptional gene regulation. Expression of this gene has been correlated with the grade of the malignancy and proliferative activity in gliomas and melanomas. A pseudogene for this gene is located on chromosome 11q13.


## References

Fan, L.F., et al. Int J Colorectal Dis 25(1):17-23(2010)  
de Sousa Abreu, R., et al. J. Biol. Chem. 284(18):12125-12135(2009)  
Murayama, M., et al. J. Gastroenterol. 44(3):173-182(2009)  
Gotte, M., et al. J. Pathol. 215(3):317-329(2008)  
Sanchez-Diaz, P.C., et al. BMC Cancer 8, 280 (2008) :

## Images




Western blot analysis of lysate from SH-SY5Y cell line, using MSI1 Antibody (N-term)(Cat. #AP11965a). AP11965a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug per lane.



MSI1 Antibody (N-term) (Cat. #AP11965a)immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining.This data demonstrates the use of MSI1 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

MSI1 Antibody (N-term) (Cat. #AP11965a) flow cytometric analysis of CEM cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated



goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.