

PRAMEF8 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP12211b

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	Q5VWM4
Other Accession	A6NMC2 , O95522 , NP_001012276.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB29974
Calculated MW	53655
Antigen Region	419-447

Additional Information

Gene ID	391002
Other Names	PRAME family member 8, PRAMEF8
Target/Specificity	This PRAMEF8 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 419-447 amino acids from the C-terminal region of human PRAMEF8.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PRAMEF8 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Name	PRAMEF8 (HGNC:24074)
Synonyms	PRAMEF24


References

Gregory, S.G., et al. Nature 441(7091):315-321(2006)


Images



PRAMEF8 Antibody (C-term) (Cat. #AP12211b) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the PRAMEF8 antibody detected the PRAMEF8 protein (arrow).



PRAMEF8 Antibody (C-term) (Cat. #AP12211b) immunohistochemistry analysis in formalin fixed and paraffin embedded human melanoma tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of PRAMEF8 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



PRAMEF8 Antibody (C-term) (Cat. #AP12211b) flow cytometric analysis of NCI-H460 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.