

# SNX13 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP12244b

## Product Information

---

Application	WB, E
Primary Accession	<a href="#">Q9Y5W8</a>
Other Accession	<a href="#">Q6PHS6</a> , <a href="#">NP_055947.1</a>
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB32232
Calculated MW	112189
Antigen Region	780-807

## Additional Information

---

Gene ID	23161
Other Names	Sorting nexin-13, RGS domain- and PHOX domain-containing protein, RGS-PX1, SNX13, KIAA0713
Target/Specificity	This SNX13 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 780-807 amino acids from the C-terminal region of human SNX13.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	SNX13 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	SNX13
Synonyms	KIAA0713

<b>Function</b>	May be involved in several stages of intracellular trafficking. May play a role in endosome homeostasis (By similarity). Acts as a GAP for Galphas.
<b>Cellular Location</b>	Early endosome membrane; Peripheral membrane protein; Cytoplasmic side


## Background

This gene encodes a PHOX domain- and RGS domain-containing protein that belongs to the sorting nexin (SNX) family and the regulator of G protein signaling (RGS) family. The PHOX domain is a phosphoinositide binding domain, and the SNX family members are involved in intracellular trafficking. The RGS family members are regulatory molecules that act as GTPase activating proteins for G alpha subunits of heterotrimeric G proteins. The RGS domain of this protein interacts with G alpha(s), accelerates its GTP hydrolysis, and attenuates G alpha(s)-mediated signaling. Overexpression of this protein delays lysosomal degradation of the epidermal growth factor receptor. Because of its bifunctional role, this protein may link heterotrimeric G protein signaling and vesicular trafficking.

## References

Rose, J. Phd, et al. Mol. Med. (2010) In press :  
Hillman, R.T., et al. Genome Biol. 5 (2), R8 (2004) :  
Worby, C.A., et al. Nat. Rev. Mol. Cell Biol. 3(12):919-931(2002)  
Kosan, C., et al. Cytogenet. Genome Res. 97 (3-4), 167-170 (2002) :  
Zheng, B., et al. Science 294(5548):1939-1942(2001)

## Images



Anti-SNX13 Antibody (C-term) at 1:1000 dilution + HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 112 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.