

# LIPC Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP12283c

## Product Information

---

<b>Application</b>	IHC-P, FC, WB, E
<b>Primary Accession</b>	<a href="#">P11150</a>
<b>Other Accession</b>	<a href="#">NP_000227.2</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB31016
<b>Calculated MW</b>	55914
<b>Antigen Region</b>	310-338

## Additional Information

---

<b>Gene ID</b>	3990
<b>Other Names</b>	Hepatic triacylglycerol lipase, HL, Hepatic lipase, Lipase member C, LIPC, HTGL
<b>Target/Specificity</b>	This LIPC antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 310-338 amino acids from the Central region of human LIPC.
<b>Dilution</b>	IHC-P~~1:100~500 FC~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	LIPC Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	LIPC
<b>Synonyms</b>	HTGL
<b>Function</b>	Catalyzes the hydrolysis of triglycerides and phospholipids present in

circulating plasma lipoproteins, including chylomicrons, intermediate density lipoproteins (IDL), low density lipoproteins (LDL) of large size and high density lipoproteins (HDL), releasing free fatty acids (FFA) and smaller lipoprotein particles (PubMed:[12032167](#), PubMed:[26193433](#), PubMed:[7592706](#), PubMed:[8798474](#)). Also exhibits lysophospholipase activity (By similarity). Can hydrolyze both neutral lipid and phospholipid substrates but shows a greater binding affinity for neutral lipid substrates than phospholipid substrates (By similarity). In native LDL, preferentially hydrolyzes the phosphatidylcholine species containing polyunsaturated fatty acids at sn-2 position (PubMed:[26193433](#)).

## Cellular Location

Secreted.


## Background

LIPC encodes hepatic triglyceride lipase, which is expressed in liver. LIPC has the dual functions of triglyceride hydrolase and ligand/bridging factor for receptor-mediated lipoprotein uptake.


## References

Reynolds, R., et al. Ophthalmology 117(10):1989-1995(2010)  
 Jablonski, K.A., et al. Diabetes 59(10):2672-2681(2010)  
 Hu, M., et al. Pharmacogenet. Genomics 20(10):634-637(2010)  
 Romero, R., et al. Am. J. Obstet. Gynecol. 203 (4), 361 (2010) :  
 Kashani Farid, M.A., et al. Lipids Health Dis 9, 96 (2010) :

## Images




Anti-LIPC Antibody (Center) at 1:500 dilution + HT-29 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 56 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



LIPC Antibody (Center) (Cat. #AP12283c) immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of LIPC Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

LIPC Antibody (Center) (Cat. #AP12283c) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated



goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.