

SNRPB2 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP12381a

Product Information

Application WB, E **Primary Accession** WB, E

Reactivity Human **Predicted** Mouse Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB31306 **Calculated MW** 25486 **Antigen Region** 1-30

Additional Information

Gene ID 6629

Other Names U2 small nuclear ribonucleoprotein B", U2 snRNP B", SNRPB2

Target/SpecificityThis SNRPB2 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 1-30 amino acids from the N-terminal

region of human SNRPB2.

Dilution WB~~1:1000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This

antibody is purified through a protein A column, followed by peptide affinity

purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions SNRPB2 Antibody (N-term) is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name SNRPB2

Function Involved in pre-mRNA splicing as component of the spliceosome

(PubMed: 11991638, PubMed: 28076346, PubMed: 28502770,

PubMed: 28781166, PubMed: 32494006). Associated with sn-RNP U2, where it

contributes to the binding of stem loop IV of U2 snRNA (PubMed: 32494006, PubMed: 9716128).

Cellular Location

Nucleus


Background

The protein encoded by this gene associates with stem loop IV of U2 small nuclear ribonucleoprotein (U2 snRNP) in the presence of snRNP-A'. The encoded protein may play a role in pre-mRNA splicing. Autoantibodies from patients with systemic lupus erythematosus frequently recognize epitopes on the encoded protein. Two transcript variants encoding the same protein have been found for this gene.

References

Oh, J.H., et al. Mamm. Genome 16(12):942-954(2005) Rush, J., et al. Nat. Biotechnol. 23(1):94-101(2005) Rush, J., et al. Nat. Biotechnol. 23(1):94-101(2005) Chung, S., et al. Biochim. Biophys. Acta 1576(3):287-297(2002) Jurica, M.S., et al. RNA 8(4):426-439(2002)

Images



All lanes: Anti-SNRPB2 Antibody (N-term) at 1:500 dilution + 293T whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 25. 5 KDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.