

SLMO2 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP12404c

Product Information

Application	WB, IHC-P, E
Primary Accession	Q9Y3B1
Other Accession	Q58DB0 , NP_057129.2
Reactivity	Human
Predicted	Bovine, Mouse, Rat, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB31423
Calculated MW	21495
Antigen Region	92-121

Additional Information

Gene ID	51012
Other Names	Protein slowmo homolog 2, SLMO2, C20orf45
Target/Specificity	This SLMO2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 92-121 amino acids from the Central region of human SLMO2.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	SLMO2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Name	PRELID3B
Synonyms	C20orf45, SLMO2


References

Deloukas, P., et al. Nature 414(6866):865-871(2001)


Images




SLMO2 Antibody (Center) (Cat. #AP12404c)immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining.This data demonstrates the use of SLMO2 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



All lanes: Anti-SLMO2 Antibody (Center) at 1:1000 dilution + Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 21 KDa Blocking/Dilution buffer: 5% NFDM/TBST.




All lanes: Anti-SLMO2 Antibody (Center) at 1:1000 dilution + Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 21 KDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes: Anti-SLMO2 Antibody (Center) at 1:1000 dilution
Lane 1: HeLa whole cell lysate Lane 2: Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 21 KDa Blocking/Dilution buffer: 5% NFDM/TBST.

All lanes: Anti-SLMO2 Antibody (Center) at 1:1000 dilution



+ Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 21 KDa Blocking/Dilution buffer: 5% NFDm/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.