

C4orf36 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP12482c

Product Information

Application WB, IHC-P, E **Primary Accession Q96KX1 Other Accession** NP 653246.1 Reactivity Human Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB27439 Calculated MW 13276 57-86 **Antigen Region**

Additional Information

Gene ID 132989

Other Names Uncharacterized protein C4orf36, C4orf36

Target/Specificity This C4orf36 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 57-86 amino acids from the Central

region of human C4orf36.

Dilution WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This

antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation

followed by dialysis against PBS.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions C4orf36 Antibody (Center) is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name C4orf36


Background

The function of this protein remains unknown.

References

Rose, J. Phd, et al. Mol. Med. (2010) In press:

Images



All lanes: Anti-C4orf36 Antibody (Center) at 1:2000 dilution Lane 1: MCF-7 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 13kDa Blocking/Dilution buffer: 5% NFDM/TBST.



C4orf36 Antibody (Center) (Cat. #AP12482c)immunohistochemistry analysis in formalin fixed and paraffin embedded human testis tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of C4orf36 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.