

WDR3 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP12525a

Product Information

Application	WB, FC, E
Primary Accession	Q9UNX4
Other Accession	NP_006775.1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB31529
Calculated MW	106099
Antigen Region	208-236

Additional Information

Gene ID	10885
Other Names	WD repeat-containing protein 3, WDR3
Target/Specificity	This WDR3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 208-236 amino acids from the N-terminal region of human WDR3.
Dilution	WB~~1:1000 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	WDR3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	WDR3 (HGNC:12755)
Function	Part of the small subunit (SSU) processome, first precursor of the small eukaryotic ribosomal subunit. During the assembly of the SSU processome in the nucleolus, many ribosome biogenesis factors, an RNA chaperone and ribosomal proteins associate with the nascent pre- rRNA and work in concert

to generate RNA folding, modifications, rearrangements and cleavage as well as targeted degradation of pre- ribosomal RNA by the RNA exosome.

Cellular Location Nucleus, nucleolus

Tissue Location Ubiquitous.


Background

This gene encodes a nuclear protein containing 10 WD repeats. WD repeats are approximately 30- to 40-amino acid domains containing several conserved residues, which usually include a trp-asp at the C-terminal end. Proteins belonging to the WD repeat family are involved in a variety of cellular processes, including cell cycle progression, signal transduction, apoptosis, and gene regulation.


References

- Akdi, A., et al. Thyroid 20(7):803-809(2010)
McMahon, M., et al. J. Biol. Chem. 285(24):18309-18318(2010)
Olsen, J.V., et al. Cell 127(3):635-648(2006)
Nousiainen, M., et al. Proc. Natl. Acad. Sci. U.S.A. 103(14):5391-5396(2006)
Nousiainen, M., et al. Proc. Natl. Acad. Sci. U.S.A. 103(14):5391-5396(2006)

Images



WDR3 Antibody (N-term) (Cat. #AP12525a) western blot analysis in HeLa, K562 cell line lysates (35ug/lane). This demonstrates the WDR3 antibody detected the WDR3 protein (arrow).



WDR3 Antibody (N-term) (Cat. #AP12525a) flow cytometric analysis of HeLa cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.