

EIF3CL Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP12609a

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	Q99613
Other Accession	B5ME19 , B5DFC8 , Q8R1B4 , Q3SYW6 , NP_001032897.1 , NP_003743.1
Reactivity	Human
Predicted	Bovine, Mouse, Rat, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB32067
Calculated MW	105344
Antigen Region	127-155

Additional Information

Gene ID	8663
Other Names	Eukaryotic translation initiation factor 3 subunit C {ECO:0000255 HAMAP-Rule:MF_03002}, eIF3c {ECO:0000255 HAMAP-Rule:MF_03002}, Eukaryotic translation initiation factor 3 subunit 8 {ECO:0000255 HAMAP-Rule:MF_03002}, eIF3 p110 {ECO:0000255 HAMAP-Rule:MF_03002}, EIF3C {ECO:0000255 HAMAP-Rule:MF_03002}
Target/Specificity	This EIF3CL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 127-155 amino acids from the N-terminal region of human EIF3CL.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	EIF3CL Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	EIF3C {ECO:0000255 HAMAP-Rule:MF_03002}
Function	Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed: 17581632 , PubMed: 25849773 , PubMed: 27462815). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl- tRNA _i and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed: 17581632). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed: 25849773).
Cellular Location	Cytoplasm {ECO:0000255 HAMAP-Rule:MF_03002}.


Background

Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA_i and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of posttermination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation.

References


Imielinski, M., et al. Nat. Genet. 41(12):1335-1340(2009)
Zhou, M., et al. Proc. Natl. Acad. Sci. U.S.A. 105(47):18139-18144(2008)
Masutani, M., et al. EMBO J. 26(14):3373-3383(2007)
Damoc, E., et al. Mol. Cell Proteomics 6(7):1135-1146(2007)
Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)

Images




All lanes: Anti-EIF3CL Antibody (N-term) at 1:1000 dilution
Lane 1: K562 whole cell lysate Lane 2: NIH/3T3 whole cell lysate Lane 3: Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution.
Observed band size: 110 KDa Blocking/Dilution buffer: 5% NFDM/TBST.

EIF3CL Antibody (N-term) (Cat. #AP12609a) immunohistochemistry analysis in formalin fixed and paraffin embedded human cerebellum tissue



followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of EIF3CL Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



EIF3CL Antibody (N-term) (Cat. #AP12609a) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.