

# KERA Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP12617b

## Product Information

---

<b>Application</b>	WB, FC, IHC-P-Leica, E
<b>Primary Accession</b>	<a href="#">O60938</a>
<b>Other Accession</b>	<a href="#">NP_008966.1</a>
<b>Reactivity</b>	Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB32135
<b>Calculated MW</b>	40509
<b>Antigen Region</b>	228-257

## Additional Information

---

<b>Gene ID</b>	11081
<b>Other Names</b>	Keratocan, KTN, Keratan sulfate proteoglycan keratocan, KERA, SLRR2B
<b>Target/Specificity</b>	This KERA antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 228-257 amino acids from the C-terminal region of human KERA.
<b>Dilution</b>	WB~~1:1000 FC~~1:25 IHC-P-Leica~~1:500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	KERA Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	KERA
<b>Synonyms</b>	SLRR2B
<b>Function</b>	May be important in developing and maintaining corneal transparency and

for the structure of the stromal matrix.

## Cellular Location

Secreted, extracellular space, extracellular matrix

## Tissue Location

Cornea (at protein level) (PubMed:10802664, PubMed:11683372). Increased expression in the stroma of keratoconus corneas (PubMed:11683372). Also detected in trachea, and in low levels, in intestine, skeletal muscle, ovary, lung and putamen (PubMed:10802664).

## Background

The protein encoded by this gene is a keratan sulfate proteoglycan that is involved in corneal transparency. Defects in this gene are a cause of autosomal recessive cornea plana 2 (CNA2).


## References

Aldave, A.J., et al. Invest. Ophthalmol. Vis. Sci. 51(8):4006-4012(2010)  
Dimasi, D.P., et al. Mol. Vis. 16, 562-569 (2010) :  
Wheeler, H.E., et al. PLoS Genet. 5 (10), E1000685 (2009) :  
Melrose, J., et al. Arthritis Res. Ther. 10 (4), R79 (2008) :  
Liskova, P., et al. Mol. Vis. 13, 1339-1347 (2007) :

## Images




Immunohistochemical analysis of paraffin-embedded human skeletal muscle tissue using AP12617b performed on the Leica® BOND RXm. Samples were incubated with primary antibody(1/500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Overlay histogram showing SK-OV-3 cells stained with AP12617b(green line). The cells were fixed with 2% paraformaldehyde and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed at 1/200 dilution for 40 min at Room temperature. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >10, 000 events was performed.

Anti-KERA Antibody (C-term) at 1:1000 dilution + Mouse eyeball lysate Lysates/proteins at 20 µg per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 41 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.