

# MyoD1 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP12646c

## Product Information

---

<b>Application</b>	IF, WB, E
<b>Primary Accession</b>	<a href="#">P15172</a>
<b>Other Accession</b>	<a href="#">NP_002469.2</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Calculated MW</b>	34501
<b>Antigen Region</b>	211-240

## Additional Information

---

<b>Gene ID</b>	4654
<b>Other Names</b>	Myoblast determination protein 1, Class C basic helix-loop-helix protein 1, bHLHC1, Myogenic factor 3, Myf-3, MYOD1, BHLHC1, MYF3, MYOD
<b>Target/Specificity</b>	This MyoD1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 211-240 amino acids from the Central region of human MyoD1.
<b>Dilution</b>	IF~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	MyoD1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	MYOD1
<b>Synonyms</b>	BHLHC1, MYF3, MYOD
<b>Function</b>	Acts as a transcriptional activator that promotes transcription of muscle-specific target genes and plays a role in muscle differentiation.

Together with MYF5 and MYOG, co-occupies muscle-specific gene promoter core region during myogenesis. Induces fibroblasts to differentiate into myoblasts. Interacts with and is inhibited by the twist protein. This interaction probably involves the basic domains of both proteins (By similarity).

#### Cellular Location

Nucleus.

## Background

---

This gene encodes a nuclear protein that belongs to the basic helix-loop-helix family of transcription factors and the myogenic factors subfamily. It regulates muscle cell differentiation by inducing cell cycle arrest, a prerequisite for myogenic initiation. The protein is also involved in muscle regeneration. It activates its own transcription which may stabilize commitment to myogenesis.


## References

---

Xynos, A., et al. Stem Cells 28(5):965-973(2010)  
Stuelsatz, P., et al. J. Biol. Chem. 285(17):12670-12683(2010)  
Hiraoka, S., et al. Hum. Pathol. 41(1):38-47(2010)  
Yerges, L.M., et al. J. Bone Miner. Res. 24(12):2039-2049(2009)  
Yang, Z., et al. Genes Dev. 23(6):694-707(2009)

## Images

---



All lanes : Anti-MyoD1 Antibody (Center) at 1:500-1000 dilution  
Lane 1: Human skeletal muscle lysate  
Lane 2: Human placenta lysate  
Lane 3: U-2OS whole cell lysate  
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 35 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.