

## PLOD3 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)  
Catalog # AP12733a

### Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">O60568</a>
<b>Other Accession</b>	<a href="#">NP_001075.1</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB32395
<b>Calculated MW</b>	84785
<b>Antigen Region</b>	78-105

### Additional Information

---

<b>Gene ID</b>	8985
<b>Other Names</b>	Procollagen-lysine, 2-oxoglutarate 5-dioxygenase 3, Lysyl hydroxylase 3, LH3, PLOD3
<b>Target/Specificity</b>	This PLOD3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 78-105 amino acids from the N-terminal region of human PLOD3.
<b>Dilution</b>	WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	PLOD3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

### Protein Information

---

<b>Name</b>	PLOD3
<b>Function</b>	Multifunctional enzyme that catalyzes a series of essential post-translational modifications on Lys residues in procollagen (PubMed: <a href="#">11956192</a> , PubMed: <a href="#">12475640</a> , PubMed: <a href="#">18298658</a> , PubMed: <a href="#">18834968</a> ,

PubMed:[30089812](#)). Plays a redundant role in catalyzing the formation of hydroxylysine residues in -Xaa-Lys-Gly- sequences in collagens (PubMed:[11956192](#), PubMed:[12475640](#), PubMed:[18298658](#), PubMed:[18834968](#), PubMed:[30089812](#), PubMed:[9582318](#), PubMed:[9724729](#)). Plays a redundant role in catalyzing the transfer of galactose onto hydroxylysine groups, giving rise to galactosyl 5-hydroxylysine (PubMed:[12475640](#), PubMed:[18298658](#), PubMed:[18834968](#), PubMed:[30089812](#)). Has an essential role by catalyzing the subsequent transfer of glucose moieties, giving rise to 1,2-glucosylgalactosyl-5-hydroxylysine residues (PubMed:[10934207](#), PubMed:[11896059](#), PubMed:[11956192](#), PubMed:[12475640](#), PubMed:[18298658](#), PubMed:[18834968](#), PubMed:[30089812](#)). Catalyzes hydroxylation and glycosylation of Lys residues in the MBL1 collagen- like domain, giving rise to hydroxylysine and 1,2-glucosylgalactosyl-5-hydroxylysine residues (PubMed:[25419660](#)). Essential for normal biosynthesis and secretion of type IV collagens (Probable) (PubMed:[18834968](#)). Essential for normal formation of basement membranes (By similarity).

#### Cellular Location

Rough endoplasmic reticulum. Endoplasmic reticulum lumen. Endoplasmic reticulum membrane {ECO:0000250 | UniProtKB:Q9R0E1}; Peripheral membrane protein {ECO:0000250 | UniProtKB:Q9R0E1}; Lumenal side {ECO:0000250 | UniProtKB:Q9R0E1}. Secreted Secreted, extracellular space {ECO:0000250 | UniProtKB:Q9R0E1}. Note=The majority of the secreted protein is associated with the extracellular matrix. {ECO:0000250 | UniProtKB:Q9R0E1}

#### Tissue Location

Ubiquitous (PubMed:9724729). Detected in heart, placenta and pancreas and at lower levels in lung, liver and skeletal muscle (PubMed:9582318, PubMed:9724729).

## Background

---

The protein encoded by this gene is a membrane-bound homodimeric enzyme that is localized to the cisternae of the rough endoplasmic reticulum. The enzyme (cofactors iron and ascorbate) catalyzes the hydroxylation of lysyl residues in collagen-like peptides. The resultant hydroxylysyl groups are attachment sites for carbohydrates in collagen and thus are critical for the stability of intermolecular crosslinks. Some patients with Ehlers-Danlos syndrome type VIB have deficiencies in lysyl hydroxylase activity.

## References


---

Wang, C., et al. J. Cell. Mol. Med. 13(3):508-521(2009)  
Salo, A.M., et al. Am. J. Hum. Genet. 83(4):495-503(2008)  
Salo, A.M., et al. J. Cell. Physiol. 207(3):644-653(2006)  
Wang, C., et al. Matrix Biol. 21(7):559-566(2002)  
Rautavaoma, K., et al. J. Biol. Chem. 277(25):23084-23091(2002)

## Images

---

PLOD3 Antibody (N-term) (Cat. #AP12733a) western blot analysis in MDA-MB435 cell line lysates (35ug/lane).This demonstrates the PLOD3 antibody detected the PLOD3 protein (arrow).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.