10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

SPOPL Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12768c

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	Q6IQ16
Other Accession	NP_001001664.1
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB26409
Calculated MW	44647
Antigen Region	221-249

Additional Information

Gene ID	339745
Other Names	Speckle-type POZ protein-like, HIB homolog 2, Roadkill homolog 2, SPOPL
Target/Specificity	This SPOPL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 221-249 amino acids from the Central region of human SPOPL.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	SPOPL Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SPOPL
Function	Component of a cullin-RING-based BCR (BTB-CUL3-RBX1) E3 ubiquitin-protein ligase complex that mediates the ubiquitination and

subsequent proteasomal degradation of target proteins, but with relatively low efficiency. Cullin-RING-based BCR (BTB-CUL3-RBX1) E3 ubiquitin-protein ligase complexes containing homodimeric SPOPL or the heterodimer formed by SPOP and SPOPL are less efficient than ubiquitin ligase complexes containing only SPOP. May function to down-regulate the activity of cullin-RING-based BCR (BTB-CUL3-RBX1) E3 ubiquitin- protein ligase complexes that contain SPOP.

Cellular Location Nucleus.


Background

In complex with a cullin, may act in ubiquitination and proteasomal degradation processes (By similarity).


References

Lamesch, P., et al. Genomics 89(3):307-315(2007)
Zhang, Q., et al. Dev. Cell 10(6):719-729(2006)
Kent, D., et al. Development 133(10):2001-2010(2006)

Images




SPOPL Antibody (Center) (Cat. #AP12768c) western blot analysis in uterus tumor cell line lysates (35ug/lane).This demonstrates the SPOPL antibody detected the SPOPL protein (arrow).



SPOPL Antibody (Center) (Cat. #AP12768c)immunohistochemistry analysis in formalin fixed and paraffin embedded human liver tissue followed by peroxidase conjugation of the secondary antibody and DAB staining.This data demonstrates the use of SPOPL Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

SPOPL Antibody (Center) (Cat. #AP12768c) flow cytometric analysis of Jurkat cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated donkey-anti-rabbit secondary antibodies were used for the analysis.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.