

# HIPK4 Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP12924a

#### **Product Information**

**Application** IHC-P, WB, E **Primary Accession Q8NE63 Other Accession** NP 653286.2 Reactivity Human Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB22867 **Calculated MW** 69425

#### **Additional Information**

**Gene ID** 147746

Other Names Homeodomain-interacting protein kinase 4, HIPK4

**Target/Specificity** This HIPK4 antibody is generated from rabbits immunized with a recombinant

protein from human HIPK4.

**Dilution** IHC-P~~1:100~500 WB~~1:1000 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** HIPK4 Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

# **Protein Information**

Name HIPK4

**Function** Protein kinase that phosphorylates human TP53 at Ser-9, and thus induces

TP53 repression of BIRC5 promoter (By similarity). May act as a corepressor

of transcription factors (Potential).

Cellular Location Cytoplasm.


# **Background**

HIPK4 is a protein kinase that phosphorylates human TP53 at Ser-9, and thus induces TP53 repression of BIRC5 promoter (By similarity). May act as a corepressor of transcription factors (Potential).


## References

Arai, S., et al. FEBS Lett. 581(29):5649-5657(2007)

### **Images**



Anti-HIPK4 Antibody at 1:1000 dilution + human testis lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 69 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



HIPK4 Antibody (Cat. #AP12924a)immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of HIPK4 Antibody for immunohistochemistry. Clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.