

LZTS1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP13069a

Product Information

Application	WB, E
Primary Accession	Q9Y250
Other Accession	NP_066300.1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	66613
Antigen Region	77-105

Additional Information

Gene ID	11178
Other Names	Leucine zipper putative tumor suppressor 1, F37/esophageal cancer-related gene-coding leucine-zipper motif, Fez1, LZTS1, FEZ1
Target/Specificity	This LZTS1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 77-105 amino acids from the N-terminal region of human LZTS1.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	LZTS1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	LZTS1
Synonyms	FEZ1
Function	Involved in the regulation of cell growth. May stabilize the active CDC2-cyclin B1 complex and thereby contribute to the regulation of the cell

cycle and the prevention of uncontrolled cell proliferation. May act as a tumor suppressor.

Cellular Location

Cytoplasm. Cell membrane. Cell projection, dendritic spine Postsynaptic density. Synapse Note=Associated with the plasma membrane and with microtubules Detected in dendritic spines, especially in the postsynaptic density (By similarity).

Tissue Location

Highly expressed in testis, prostate, spleen, thymus, ovary and brain. Detected at lower levels in heart, placenta, small intestine, colon, liver, kidney, skeletal muscle and pancreas Not detectable in primary tumors from breast and prostate and in many cancer cell lines.


Background

This gene encodes a tumor suppressor protein that is ubiquitously expressed in normal tissues. In uveal melanomas, expression of this protein is silenced in rapidly metastasizing and metastatic tumor cells but has normal expression in slowly metastasizing or nonmetastasizing tumor cells. This protein may have a role in cell-cycle control by interacting with the Cdk1/cyclinB1 complex. This gene is located on chromosomal region 8p22. Loss of heterozygosity (LOH) in the 8p arm is a common characteristic of many types of cancer.

References

Olson, J.E., et al. Breast Cancer Res. Treat. (2010) In press :
Chattopadhyay, I., et al. Mutat. Res. 696(2):130-138(2010)
Califano, D., et al. J. Cell. Physiol. 222(2):382-386(2010)
Shyn, S.I., et al. Mol. Psychiatry (2009) In press :
Canova, C., et al. Cancer Res. 69(7):2956-2965(2009)

Images



LZTS1 Antibody (N-term) (Cat. #AP13069a) western blot analysis in A2058 cell line and mouse brain tissue lysates (35ug/lane). This demonstrates the LZTS1 antibody detected the LZTS1 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.