

PQLC2 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP13107b

Product Information

Application Primary Accession	WB, IHC-P, E <u>Q6ZP29</u>
Other Accession	<u>NP_060235.2</u> , <u>NP_001035214.1</u>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB31429
Calculated MW	31947
Antigen Region	222-251

Additional Information

Gene ID	54896
Other Names	Lysosomal amino acid transporter 1 homolog, PQ-loop repeat-containing protein 2, PQLC2
Target/Specificity	This PQLC2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 222-251 amino acids from the C-terminal region of human PQLC2.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PQLC2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SLC66A1 (<u>HGNC:26001</u>)
Function	Amino acid transporter that specifically mediates the pH- dependent export of the cationic amino acids arginine, histidine and lysine from lysosomes.

Background

The specific function of this protein remains unknown. There are 3 named isoforms.

References

Gregory, S.G., et al. Nature 441(7091):315-321(2006)

Images



All lanes: Anti-PQLC2 Antibody (C-term) at 1:1000 dilution + Human crebellum lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 36 KDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.