

PLS1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP13147b

Product Information

Application	WB, IHC-P, E
Primary Accession	<u>Q14651</u>
Other Accession	<u>NP_001165783.1</u> , <u>NP_001138791.1</u>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB32565
Calculated MW	70253
Antigen Region	550-579

Additional Information

Gene ID	5357
Other Names	Plastin-1, Intestine-specific plastin, I-plastin, PLS1
Target/Specificity	This PLS1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 550-579 amino acids from the C-terminal region of human PLS1.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PLS1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PLS1
Function	Actin-bundling protein. In the inner ear, it is required for stereocilia formation. Mediates liquid packing of actin filaments that is necessary for stereocilia to grow to their proper dimensions.

Cellular Location	Cytoplasm {ECO:0000250 UniProtKB:Q3V0K9}. Cell projection, stereocilium {ECO:0000250 UniProtKB:Q3V0K9}
Tissue Location	In small intestine, colon, and kidney; relatively lower levels of expression are detected in the lung and stomach

Background

Plastins are a family of actin-binding proteins that are conserved throughout eukaryote evolution and expressed in most tissues of higher eukaryotes. In humans, two ubiquitous plastin isoforms (L and T) have been identified. The protein encoded by this gene is a third distinct plastin isoform, which is specifically expressed at high levels in the small intestine. Alternatively spliced transcript variants varying in the 5' UTR, but encoding the same protein, have been found for this gene. A pseudogene of this gene is found on chromosome 11.

References

Chafel, M.M., et al. Dev. Dyn. 203(2):141-151(1995) Lin, C.S., et al. Mol. Cell. Biol. 14(4):2457-2467(1994) Shibata, M., et al. J. Leukoc. Biol. 54(1):10-16(1993) Lin, C.S., et al. J. Biol. Chem. 268(4):2781-2792(1993) Zu, Y., et al. Biochemistry 29(4):1055-1062(1990)

Images



PLS1 Antibody (C-term) (Cat. #AP13147b) western blot analysis in Jurkat cell line lysates (35ug/lane).This demonstrates the PLS1 antibody detected the PLS1 protein (arrow).



PLS1 Antibody (C-term) (Cat.

#AP13147b)immunohistochemistry analysis in formalin fixed and paraffin embedded human kidney tissue followed by peroxidase conjugation of the secondary antibody and DAB staining.This data demonstrates the use of PLS1 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

Citations

• Enteral delivery of proteins enhances the expression of proteins involved in the cytoskeleton and protein biosynthesis in human duodenal mucosa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.